

ICONES PLANTARUM;

FIGURES,

BRIEF DESCRIPTIVE CHARACTERS AND REMARKS

OF

NEW OR RARE PLANTS,

SELECTED FROM THE AUTHOR'S HERBARIUM.

BY SIR WILLIAM JACKSON HOOKER, K.H.,

L.L.D., F.R.A.; AND L.S.

MEMBER OF THE IMP. ACAD. NAT. CURE, ETC., ETC., ETC.

HONORARY MEMBER OF THE ROYAL IRISH ACADEMY, OF THE ROYAL MEDICAL AND CHIRURGICAL SOC. OF LONDON, ETC., ETC.

AND

DIRECTOR OF THE ROYAL BOTANIC GARDEN, K.W.

VOL. IV.

LONDON:

LONGMAN, ORME, BROWN, GREEN, AND LONGMANS.

M D C C C X L I.

LONDON :
Printed by A. SPOTTISWOODE,
New-Street-Square.

INDEX

TO THE

PLANTS CONTAINED IN VOLUME IV.;

ARRANGED ACCORDING TO THE NATURAL ORDERS.

	TAB.		TAB.
PAPAVERACEÆ.		PASSIFLOREÆ.	
Meconella Oregana, Nutt.	360	Passiflora Leschenaultii, De Cand.	359
FUMARIACEÆ.		MALVACEÆ.	
Fumaria micrantha, Lag.	363	Sida parnassiaefolia, Hook.	385
ANONACEÆ.		TERNSTEINMIACEÆ.	
Anona bibracteata, Hook.	328	Saurauja barbigeria, Hook.	331
CRUCIFERÆ.		pedunculata, Hook.	341, 342
Arabis puberula, Nutt.	359	EUPHORBIACEÆ.	
SAPINDACEÆ.		Mozinna spathulata, Ort.	357
Fappea capensis, Eckl. et Zeyh.	352	Euphorbia sphaerorhiza, Benth.	346
DROSERACEÆ.		CELASTRINÆ.	
Drosera bulbosa, Hook.	375	Wimmeria concolor, Chom. et Schlech.	353
macrophylla, Lind.	376	LEGUMINOSÆ.	
stolonifera, Endl.	389	Crotalaria bupleurifolia, Schlecht.	372
BERBERIDÆ.		Mimosa floribunda, H. B. K.	373
Berberis (Mahonia) fraxinifolia, Hook.	329, 390	Acacia setigera, Hook.	316
OLACINÆ.		squamata, Lindl.	367
Ximenia parviflora, Benth.	350	diptera, Lindl.	369
OSALIDÆ.		incrensata, Zude.	370
Oxalis Cratensis, Hook.	361	ROSACEÆ.	
Rubus roseifolius, Sm.		Rubus roseifolius, Sm.	349
Prunus (Cerasus) samyoides, Chom. et Schlech.		Prunus (Cerasus) samyoides, Chom. et Schlech.	371

	TAB.		TAB.
Cercocarpus betulæfolius, Nutt.	322	VACCINIEÆ.	
parvifolius, Nutt.	323	Vaccinium Forbesii, Hook.	345
ledifolius, Nutt.	324		
		ERICÆ.	
RHEOPHOREÆ.		Gaultheria tomentosa, H. B. K.	393
Kandelia Rhedei, Wight et Arn.	362		
Bruguiera Rhedei, Arn.	397		
Malabarica, Arn.	398		
		PROTEACEÆ.	
ONAGRARIÆ.		Anadenia Mangelsii, Griseb.	337
Oenothera (Holostigma) alyssoides,			
<i>H. et Arn.</i> 339			
graciliflora,			
<i>H. et Arn.</i> 338			
		SCROPHULARIINEÆ.	
HALORAGÆ.		Harveya purpurea, Harv.	351
Meionectes Brownii, J. D. Hook.	306	Aulaya capensis, Harv.	400
Goniocarpus vernicosus, J. D. Hook.	311		
		LOBELIACEÆ.	
MYRTACEÆ.		Lobelia trigona, Rox.	358
Leptospermum rupestre, J. D. Hook.	308		
Bæckia micrantha, De Cand.	309		
		GESNERIACEÆ.	
CRASSULACEÆ.		Gloxinia Sarmentiana, Gard.	378
Tillæa macrantha, J. D. Hook.	310		
		HYDROPHYLLÆ.	
CUNONIACEÆ.		Eutoca arietoides, Hook. et Arn.	355
Weinmannia biglandulosa, A. Cunn.	301	Eutoca? lutea, Hook. et Arn.	354
		APOCYNEÆ.	
UMBELLIFERÆ.		Rauwolfia heterophylla, Willd.	366
Hydrocotyle tripartita, Br.	312		
cordifolia, J. D. Hook.	303		
Didiscus humilis, J. D. Hook.	304		
pilosus, Hugel	307		
Xanthosia dissecta, J. D. Hook.	302		
Petroselinum prostratum, De Cand.	305		
		STRYCHNÆ.	
LOANTHÆ.		Strychnos toxifera, Schomb.	364, 365
Loranthus Schiedeanus, Schlecht.	374		
Viscum falcatum, Cham. et Schlecht.	368		
		SOLANÆ.	
COMPOSITE.		Fabiana imbricata, Ruiz et Pav.	340
Actinolepis multicaulis, De Cand.	325	Cestrum vestitum, Hook.	381
Hartmannia? pungens, Hook. et Arn.	334		
		GARRYACEÆ.	
Madaraglossa heterotricha, De Cand.	326	Garrya Fadyenii, Hook.	333
Monolopia major, De Cand.	344		
minor, De Cand.	349		
Cotula myriophyloides, Harv.	335		
Helichrysum Gunnianum, Hook.	320		
Helipterum incanum, De Cand.	318		
		URTICÆ.	
Dorstenia asaroides, Gard.	399		
		PIPERACEÆ.	
Peperomia Galeottiana, Hook.	327		
lancifolia, Hook.	392		
		CUPULIFERÆ.	
Quercus agrifolia, Née	377		
densiflora, Hook. et Arn.	380		
Douglasii, Hook. et Arn.	382, 383		
		TAXINÆ.	
Taxodium sempervirens, Lamb.?	379		

INDEX.

V

	TAB.		TAB.
	CYTINÆ.		POLYPODIUM.
Cytinus dioicus, <i>Juss.</i> . . .	336	Polyodium (Eupolyp.) Hartwegi-	
		anum, <i>Hook.</i> 390	
		subpetiolatum,	
		<i>Hook.</i> 391, 392	
	ORCHIDÆ.		piligerum,
Epidendrum rigidum, <i>Jacq.</i> . . .	314	<i>Hook.</i> 321	
<i>microbulbon</i> , <i>Hook.</i> . . .	347	Allosorus Karwinskii, <i>Kunze</i> 387, 388	
Maxillaria nana, <i>Hook.</i> . . .	315	Antrophyum ensiforme, <i>Hook.</i> 394	
<i>angustifolia</i> , <i>Hook.</i> . . .	348	Aspidium (Nephrolepis) davallio-	
<i>curtipes</i> , <i>Hook.</i> . . .	384	ides, <i>Sie.</i> 395, 396	
Scaphyglottis rosea, <i>Hook.</i> . . .	313		
<i>fasciculata</i> , <i>Hook.</i> . . .	317		
	FILICES.		MUSCI.
Polypodium (Eupolyp.) discolor, <i>Hook.</i>	386	Dicranum macrodon, <i>Hook.</i> . . .	319

INDEX

TO THE

PLANTS CONTAINED IN VOLUME IV.;

ALPHABETICALLY ARRANGED.

	TAB.		TAB.
<i>Acacia diptera</i> , <i>Lindl.</i>	369	<i>Drosera bulbosa</i> , <i>Hook.</i>	375
<i>incrassata</i> , <i>Hook.</i>	370	<i>macrophylla</i> , <i>Lindl.</i>	376
<i>setigera</i> , <i>Hook.</i>	316	<i>stolonifera</i> , <i>Endl.</i>	389
<i>squamata</i> , <i>Lindl.</i>	367	<i>Epidendrum microbulbon</i> , <i>Hook.</i>	347
<i>Actinolepis multicaulis</i> , <i>De Cand.</i>	325	<i>rigidum</i> , <i>Jacq.</i>	314
<i>Allosorus Karwinskii</i> , <i>Kunze</i>	387	<i>Euphorbia sphaerorhiza</i> , <i>Benth.</i>	346
<i>Anadenia Manglesii</i> , <i>Grah.</i>	337	<i>Eutoca arietoides</i> , <i>Hook. et Arn.</i>	355
<i>Anona libracteata</i> , <i>Hook.</i>	328	<i>lutea</i> , <i>Hook. et Arn.</i>	354
<i>Antrophyum ensiforme</i> , <i>Hook.</i>	394	<i>Fabiana imbricata</i> , <i>Ruiz et Pav.</i>	340
<i>Arabis puberula</i> , <i>Nutt.</i>	359	<i>Fumaria micrantha</i> , <i>Lag.</i>	363
<i>Aspidium (Nephrolepis) davallio-</i> <i>ides</i> , <i>Sw.</i>	395, 396	<i>Garrya Fadyenii</i> , <i>Hook.</i>	333
<i>Aulaya capensis</i> , <i>Hare.</i>	400	<i>Gaultheria tomentosa</i> , <i>H. B. K.</i>	393
<i>Beckia micrantha</i> , <i>De Cand.</i>	309	<i>Gloxinia Sarmentiana</i> , <i>Gard.</i>	378
<i>Berberis (Mahonia) fraxinifolia</i> , <i>Hook.</i>	329, 330	<i>Goniocarpus vernieotus</i> , <i>J. D. Hook.</i>	311
<i>Bruguiera malabarica</i> , <i>Arn.</i>	398	<i>Hartmannia? pungens</i> , <i>Hook. et Arn.</i>	334
<i>Rheedei</i> , <i>Arn.</i>	397	<i>Harveya purpurea</i> , <i>Hare.</i>	351
<i>Cercocarpus betulafolius</i> , <i>Nutt.</i>	322	<i>Helichrysum Gunnianum</i> , <i>Hook.</i>	320
<i>ledifolius</i> , <i>Nutt.</i>	324	<i>Heptipetalum incanum</i> , <i>De Cand.</i>	318
<i>parvifolius</i> , <i>Nutt.</i>	323	<i>Hydrocotyle cordifolia</i> , <i>J. D. Hook.</i>	303
<i>Cestrum vestitum</i> , <i>Hook.</i>	381	<i>tripartita</i> , <i>Br.</i>	312
<i>Cotula myriophyloides</i> , <i>Hare.</i>	335	<i>Kandelia Rheedei</i> , <i>Wight et Arn.</i>	362
<i>Crotalaria bipinnatifolia</i> , <i>Schlecht.</i>	372	<i>Leptospermum rupestre</i> , <i>J. D. Hook.</i>	308
<i>Cytinus dioicus</i> , <i>Juss.</i>	336	<i>Lobelia trigona</i> , <i>Rox.</i>	358
<i>Dieramum macroodon</i> , <i>Hook.</i>	319	<i>Loranthus Schiedeanus</i> , <i>Schlecht.</i>	374
<i>Didesmus humilis</i> , <i>J. D. Hook.</i>	304	<i>Madaraglossa heterotricha</i> , <i>De Cand.</i>	326
<i>pilosus</i> , <i>Hegel.</i>	307	<i>Maxillaria angustifolia</i> , <i>Hook.</i>	348
<i>Dorstenia sessilifolia</i> , <i>Gard.</i>	399	<i>curtipes</i> , <i>Hook.</i>	384
		<i>nana</i> , <i>Hook.</i>	315
		<i>Meconella Oregana</i> , <i>Nutt.</i>	360

TAB.		TAB.	
Meionectes Brownii, <i>J. D. Hook.</i>	306	Prunus (<i>Cerasus</i>) samydeoides, <i>Cham.</i>	
<i>Mimosa floribunda</i> , <i>H. B. K.</i>	373	et <i>Schlecht.</i>	371
Monolopia major, <i>De Cand.</i>	344	<i>Quercus agrifolia</i> , <i>Née</i>	377
<i>minor</i> , <i>De Cand.</i>	343	<i>densiformis</i> , <i>Hook. et Arn.</i>	380
<i>Mosinna spathulata</i> , <i>Ort.</i>	357	<i>Douglasii</i> , <i>Hook. et Arn.</i>	
<i>Onothea</i> (<i>Holostigma</i>) <i>alyssoides</i> ,			382, 383
<i>Hook. et Arn.</i>	339	<i>Rauwolfia heterophylla</i> , <i>Willd.</i>	366
<i>graciliflora</i> , <i>Hook.</i>		<i>Rubus roseifolius</i> , <i>Sm.</i>	349
<i>et Arn.</i>	338	<i>Saurauja barbigera</i> , <i>Hook.</i>	331
<i>Oxalis Cratensis</i> , <i>Hook.</i>	361	<i>pedunculata</i> , <i>Hook.</i>	341, 342
<i>Pappea capensis</i> , <i>Echl. et Zeyh.</i>	352	<i>Scaphyglottis</i> ? <i>fasciculata</i> , <i>Hook.</i>	317
<i>Passiflora Leschenaultii</i> , <i>De Cand.</i>	353	<i>rosea</i> , <i>Hook.</i>	313
<i>Peperomia Galeottiana</i> , <i>Hook.</i>	328	<i>Sida parnassiaefolia</i> , <i>Hook.</i>	384
<i>lancifolia</i> , <i>Hook.</i>	332	<i>Strychnos toxifera</i> , <i>Schomb.</i>	364, 365
<i>Petroselinum prostratum</i> , <i>De Cand.</i>	305	<i>Taxodium sempervirens</i> , <i>Lamb.</i> ?	379
<i>Polypodium</i> (<i>Eupolyp.</i>) <i>discolor</i> ,		<i>Tilia macrantha</i> , <i>J. D. Hook.</i>	310
<i>Hook.</i>	386	<i>Vaccinium Forbesii</i> , <i>Hook.</i>	345
<i>Hartwegianum</i> ,		<i>Viscum falcatum</i> , <i>Cham. et Schlecht.</i>	368
<i>Hook.</i>	390	<i>Weinmannia biglandulosa</i> , <i>A. Cunn.</i>	301
<i>piligerum</i> , <i>Hook.</i>	321	<i>Wimmeria concolor</i> , <i>Cham. et</i>	
<i>subpetiolatum</i> ,		<i>Schlecht.</i>	356
<i>Hook.</i>	391, 392	<i>Xanthosia dissecta</i> , <i>J. D. Hook.</i>	303
		<i>Ximenia parviflora</i> , <i>Benth.</i>	350



TAB. CCCI.

WEINMANNIA BIGLANDULOSA.

Folis simplicibus elliptico-lanceolatis obtusis grosse obtuse serratis in petiolum brevem articulatis ad articulum minute biglandulosis, pedunculis axillaribus solitariis unifloris petiolum subæquantibus bibracteatis, antheris longe acuminatis.

Weinmannia biglandulosa. *All. Cunn. Mst. in Herb. Nostr.*

HAB. Van Dieman's Land, where it is called "Native Beech;" *All. Cunningham, Esq.* Particularly abundant on the banks of the Emu River, in the Hampshire hills. *Ronald Gunn, Esq.* *Mr Lawrence.*

Mr Cunningham observes that this shrub, or tree, grows to the "height of 30 or 35 feet." Mr Gunn's remarks are, "It is most abundant about Emu River, forming almost impenetrable scrubs (as we call thickets in Van Dieman's Land). The stems are flexuose, bending in and out in all directions with one another; and, being pretty tough, our friends James Backhouse and Dr Milligan called the plants 'right and left scrub;' as they had to work to the right and left to get out of them. The stems, which lie in all directions, are from 8 to 18 feet long, or thereabouts, but seldom exceeding 8 feet in perpendicular height. Of course situation affects it a good deal. Soil, rich alluvial."—I have hesitated whether or not there were characters sufficient to constitute a genus of this plant:—but, except the 2-seeded cells (if that number be constant) of the capsule, the much acuminate anthers, the very narrow petals, the broad and flat hypogynous disk, and the very different inflorescence, it seems sufficiently to accord with the simple-leaved *Weinmannæ*.

Fig. 1. Flower and bracts. *f. 2.* Flower laid open. *f. 3.* Petal. *f. 4.* Stamen. *f. 5.* Anther. *f. 6.* Pistil. *f. 7.* Ovary laid open:—*magnified.*



Gunnianæ.

N. O. Umbelliferae.

TAB. CCCII.

XANTHOSIA DISSECTA. J. D. Hook.

*Pusilla prostrata glabriuscula, foliis longe petiolatis tripartitis
laciiniis dissectis, umbellis pedunculatis axillaribus terminali-
busque, sepalis ovatis acuminatis, petalis lanceolatis longe
acuminatis inflexis, fructu late cordato, mericariis jugis 5
elevatis, secundariis similibus.*


HAB. Circular Head and Rocky Cape, Van Dieman's Land.

R. Gunn, Esq. (n. 880).

A small, much branched, slender, wiry annual. Stem short, branching from the base, and there very leafy. Leaves small, tripartite, and much cut and divided. Umbels small, 2-5-flowered. Involucre of 2-3 leaves. Calyx-lobes or sepals erect. Petals narrow, entire, reddish. Styles curved. Stylopodium large and conspicuous. Fruit broadly cordate, compressed.

The habit of this is very different from any described *Xanthosia*; and should perhaps, with one or two others in Herb. Hook., form a separate section of the Genus. The carpels exactly resemble those of the sections *Leucolæna* and *Euzanthosia*. *J. D. H.*

Fig. 1. Flower. *f. 2.* The same more expanded. *f. 3.* Petal. *f. 4, 5.* Fruits. *f. 6.* Vertical section of a mericarp. *f. 7.* Transverse section:—*magnified.*



Gunnianæ.

N. O. Umbelliferæ.

TAB. CCCIII.

HYDROCOTYLE CORDIFOLIA. J. D. Hook.

Subcaulis, foliis longe petiolatis cordatis glabris marginibus obscure sinuatis apice rotundatis, umbellis subbifloris longe pedunculatis, floribus majusculis, involucris diphyllois, calycis laciniis ovatis acutis, fructibus pilosis.

HAB. Common in marshy places on sandy soils, Van Dieman's Land. *R. Gunn, Esq.* (*n.* 876).

Radix e fibris paucis crassiusculis. *Folia* pleraque radicalia longe petiolata, cordata, obtusa, glabra, basi profunde biloba. *Petiolus* subhirsutus, purpurascens. *Pedunculi* longitudine petiolorum, nunc elongati, prolixi et foliosi: foliis parvis purpurascens. *Involucrum* diphyllo, foliolis concavis. *Umbella* subbiflora, floribus majusculis. *Ovarium* cordatum, pilosum. *Calyx* e laciniis deciduis vel foliolis 5 majusculis, ovatis, acutis vel acuminatis, erectis. *Petala* ovato-lanceolata, erecta, apice acuta, inflexa. *Petala* calyce longiora. *Fructus* lato-cordatus compressus, superne hirsutus. *J. D. H.*

Fig. 1. Flower. *f. 2.* Fruit. *f. 3.* Transverse section of the same:—*magnified.*



Gunnianæ.

N. O. Umbelliferæ.

TAB. CCCIV.

DIDISCUS HUMILIS. J. D. Hook.

Acaulis, foliis longe petiolatis cordato-ovatis 3-5-lobis basi connatis, subtus petiolis scapisque (folia superantibus demum prostratis) patentí-pilosis, umbellis simplicibus, petalis obovatis patentibus, fructibus biscutatis subrugosis disco depresso marginato.

HAB. Moist open plains on the Hampshire hills, Van Dieman's Land. *R. Gunn, Esq.* (n. 245.)

Radix fusiformis. *Folia omnia radicalia, numerosa, longe petiolata.* *Scapi* folio duplo longiores, demum fructiferi, decumbentes. *Umbella* simplex, multiflora, floribus parvis. *Involucrum* polyphyllum: foliolis linear-i-angustis, umbellam aequantibus. *Pedicelli* flore longiores. *Dentes calycinae* obsoletæ. *Petala* majuscula, patentia. *Fructus* compressus, biscutatus, rugosus.

The very old scapes are always seen to be bent down in a very remarkable manner, as if they had lain prostrate on the ground: the umbel of fruit curving upwards at its apex. *J. D. H.*

Fig. 1. Flower. *f. 2.* Petal. *f. 3.* Unripe fruit. *f. 4.* Ripe fruit:—*magnified.*



TAB. CCCV.

PETROSELINUM PROSTRATUM.

Caule procumbente flexuoso, foliis pinnatisectis segmentis petiolatis pinnatifidis lacinijis 5 lanceolatis ultimis apice trifidis, umbellis subsessilibus oppositifoliis, involucro submonophylo.
DC.

Petroselinum prostratum. *De Cand. Prodr.* 4. p. 102.

Apium prostratum. *Labill. Nov. Holl.* p. 76. t. 103.

β. foliorum segmentis angustioribus. Apium prostratum. *Vent. Malm.* t. 81. (TAB. NOSTRA. CCCV.)

HAB. Sands and marshes near salt or brackish water, Van Dieman's Land. *R. Gunn, Esq.* (n. 386, 68). Port Arthur.
Mr Backhouse.

There are two appearances of this plant, which at first sight are strikingly different:—the one is that figured by Labillardière, with broad segments to the leaves; the other, that represented by Ventenat and by us, with narrow segments. Both kinds are sent by Mr Gunn; and they seem to be not uncommon in various parts of the island where there are salt or brackish soils.

Fig. 1. Flower. *f. 2.* Petal. *f. 3.* Immature fruit. *f. 4.* Ripè fruit. *f. 5.* Section of a carpel, showing the jugæ and the position of the vittæ:—*magnified.*



TAB. CCCVI.

MEIONECTES BROWNII. J. D. Hook.

Meionectes Brownii.

HAB. Van Dieman's Land, near Circular Head, in wet places.

Mr R. Gunn, (n. 883.)

Herba inconspicua, aquatica, glaberrima. *Caulis* debiles, ramosi, basi radicantes, spithamei, tenuis. *Folia* alterna, pectinata, v. laciniate, subteretia; inferiora capillaceo-multifida, glaberrima, crassiuscula, opaca, lurido-virescentia, segmentis acutis, 1-nerviis; juniora minus dissecta. *Pedunculi* axillares, solitarii, $\frac{1}{2}$ lin. longi, basi bibracteolati. *Flores* minimi, inconspicui, $\frac{1}{2}$ lin. longi, penduli. *Calycis* *tubus* urceolatus, ovarium includens, eoque adhaerens, compressus, bisulcatus, limbo bipartito, lacinias oppositis late ovatis obtusis trinerviis. *Petala* 2, obovata, cymbiformia, ad apicem valde rotundata, dentibus calycis longiora, opposita, æqualia, enervia. *Stamina* 4, 2 petalis opposita, 2 iis alterna, filamentis brevibus. *Antheræ* oblongæ, subquadratae, biloculares, flavæ, longitudin-aliter dehiscentes. *Styli* 2, breves, conico-subulati, basi coaliti, hirtelli. *Ovarium* ovatum, compressum, bisulcatum, bilocular, 2-ovulatum. *Ovula* pendula. *Fructus* nucumentaceus, 2-3 lin. longus, ovatus, compressus, rugulosus, cum dentibus persistentibus calycis coronatus, 2-locularis, 2-spermus. *Semina* alba, oblongo-elliptica. *Embryo* rectus, centralis, in albumine carnosus.

A most distinct and remarkable genus of Mr Brown, who mentions it in the botanical appendix to Flinders' voyage, as containing only half the number of parts to the flower than the other genera of *Halorageæ* possess. Whether this be the species on which he founded the genus or not, I am unable to say. The ripe seed-vessels are large in proportion to the size of the flower, which is very diminutive, and hangs down from the axils of the leaves. *J. D. H.*

Fig. 1. Flowers. *f. 2.* The same, with the petals removed. *f. 3.* Young ovary. *f. 4.* Ripe fruit. *f. 5.* The same cut transversely; and *f. 6.*, cut longitudinally:—*magnified*.



Gunnianæ.

N. O. Umbelliferæ.

TAB. CCCVII.

DIDISCUS PILOSUS.

Foliis inferioribus longe-petiolatis palmatisectis segmentis cuneatis incisis dentatisque petiolisque piloso-hispidis, caulinis paucis parvis, pedunculis elongatis ramisque glabris. *Hugel.*

Didiscus pilosus. Hugel, Enum. Pl. Nov. Holl. p. 54, (in note.)

HAB. Dry sandstone hills north of Bathurst. *All. Cunningham, Esq.* Circular Head, Van Dieman's Land. *R. Gunn, Esq.* (n. 825.)

Mr Gunn finds this plant growing in poor sandy land between Circular Head and Woolnorth, but none of it east from Circular Head to Emu Bay. It is $1\frac{1}{2}$ to 2 feet high. Root fusiform. Petioles hairy and the leaves also, but on the latter the hairs are deciduous. Segments of the caudine leaves linear. Involucre of several narrow, linear leaves. Petals large, obovate, entire. Filaments of the stamens long. Fruit scabrous, of 2 orbicular mericarps, exhibiting no evident jugæ except that one which borders an elevated disk on each side the commissure. *J. D. H.*

Fig. 1. Flower. f. 2. Petal. f. 3. Fruit. f. 4. Transverse section of the fruit:—magnified.



TAB. CCCVIII.

LEPTOSPERMUM RUPESTRE. J. D. Hook.

Ramosum prostratum humile, foliis parvis obovatis ellipticisve brevissime petiolatis apice obtusis punctatis nitidis glaberrimis obscure 1-nerviis, floribus subsessilibus, calycibus hemisphaericis sub lente puberulis, limbo glabro deciduo colorato.

HAB. Rocks on the summit of Mount Wellington and other mountains in Van Dieman's Land, at an elevation of 3000 feet above the level of the sea. *Mr Lawrence. Fraser. R. Gunn, Esq. (n. 295.)* New Zealand. *Fraser in Herb. Hook.*

A small, prostrate, much branched shrub. Branches 2-3 feet long, spreading, their apices downy. Leaves numerous, alternate, coriaceous, obovate-elliptical, scarcely petiolate, 4-6 lines long, entire, smooth, of a yellowish or brownish-green colour. Flowers numerous, solitary in the axils of the leaves, on very short peduncles. Calyces turbinate, the tube, in the buds, especially downy, the teeth broad, spreading, obtuse, membranaceous, deciduous, red in the centre, with brown, diaphanous margins. Petals whitish. Stamens numerous. Fruit of 5 cells, surrounded by the semi-superior calyx, from the tube of which the epidermis readily peels off. *J. D. H.*

Fig. 1. Flower. *f. 2.* Germen. *f. 3.* Do. cut open. *f. 4.* Ripe fruit. *f. 5.* The same cut open longitudinally. *f. 6 and 7.* Seeds:—*magnified.*



TAB. CCCIX.

BÆCKEA MICRANTHA. J. D. Hook.

Folis obovato-triquetris obtusis quadrifariam imbricatis grosse punctatis glabris, floribus axillaribus solitariis subsessilibus.
 α . erecta, major. B. micrantha. *De Cand. Prodr. v. 3. p. 230.*
 β . humilis, procumbens.

HAB. The mountainous region about May-day Plain and the Black Bluff and Vale of Belvoir, Van Dieman's Land. *R. Gunn, Esq. (n. 306; α .—n. 815; β .) Mr Lawrence.*

A much branched shrub, growing in mountainous regions, erect and 3 to 7 feet high in α ; humble and prostrate in β . Branches covered with the scars of the old foliage. Leaves closely imbricated, blunt, channelled above, rounded and keeled beneath, patent, crowded, glabrous, shortly petiolated, shining, with several large round glands or dots on each face, 2-3 lines long, dark green, often tipped with red. Peduncles straight, much shorter than the leaf. Flowers small. Calyx turbinate, smooth, the limb of 5 teeth, coloured and blunt at the point. Stamens 10, short, incurved, glabrous. Fruit shortly pedunculated, of a light brown colour, somewhat elongated, smooth, crowned by the five persistent teeth of the calyx. *J. D. H.*

Fig. 1. Leaves. *f. 2, 3.* Flowers. *f. 4.* Petal. *f. 5.* Ripe fruit:—*magnified.*



Gunnianæ.

N. O. Crassulaceæ.

TAB. CCCX.

TILLÆA MACRANTHA. J. D. Hook.

Caulibus di-trichotomis, foliis oblongo-cylindraceis acutis, floribus terminalibus tetrameris, sepalis petala superantibus ciliatis glandulis linearibus nigris obsitis.

HAB. Van Dieman's Land. *R. Gunn, Esq.*

A small annual plant growing in tufts, probably in moist places. Stems much branched in a di- or trichotomous manner, scarcely more than 2 inches high. Leaves opposite, oblongo-cylindrical, acute, fleshy. Flowers terminal, solitary, on the ultimate branches or peduncles. Calyx of 4, nearly erect, elliptical-oblong, concave, acute, ciliated sepals, streaked, both within and without, with numerous black linear glands or short lines. Petals 4, ovate, slightly combined at their base, shorter than the calyx. Stamens 4, alternating with the petals. Ovaries 4, alternating with the stamens. Fruit of 4, erect, or slightly spreading, oval-cylindrical follicles, with a short style and obtuse stigma. Seeds oblongo-reniform, dotted. *J. D. H.*

Fig. 1. Flower. f. 2. The same spread open. f. 3. Petals and stamen. f. 4. Follicles. f. 5. Single follicle. f. 6. Seed: —magnified.



Gunnianæ.

N. O. Halorageæ.

TAB. CCCXI.


GONIOPARPUS VERNICOSUS. J. D. Hook.

Nanus minute pubescenti-scaber, foliis oppositis ovatis marginatis integerrimis superne vernicosis, floribus solitariis axillaribus sessilibus basi bibracteatis, petalis cymbiformibus acuminatis extus hirsutis.

HAB. This was selected from other Van Dieman's Land plants with which it was found mixed.

Allied to *Goniocarpus serpyllifolius*, (J. D. Hook. in *Ic. Pl. t. 210*); but a much smaller plant, more compact, with crowded and perfectly entire leaves, quite glossy and as if varnished on the upper-side (whence the specific name). The flowers also are similar, but the stigmas are much less feathery, and the stamens are small, (abortive?)

Fig. 1. Leaf and flower. *f. 2.* Flower expanded. *f. 3.* Petal. *f. 4.* Stamen. *f. 5.* Stigma. *f. 6.* Leaf and bracteas;—*magnified.*



Gunnianæ.

N. O. Umbelliferæ.

TAB. CCCXII.

HYDROCOTYLE TRIPARTITA.


Pilosula, foliis palmatisectis, segmentis 3 cuneatis inciso-dentatis lateralibus saepe bifidis, pedunculis petiolo multo brevioribus (nunc longioribus), umbella subcapitata 6-10-flora, fructibus orbiculato-didymis utrinque bicostatis. *DC.*

H. tripartita. *Brown in Rich. Hydr.* n. 46. f. 25. *Sieb. Exsicc.* n. 628. *De Cand. Prodr.* v. 4. p. 65.

Van Dieman's Land. *Mr Gunn.* (n. 547.)

Mr Gunn's specimens are smaller and less hairy than those figured in Richard's Monograph, but they vary in size. The two lateral ridges on the carpels are united into one; between them and the commissure there is, on each side, a small deep sunken pit or hollow, which gives to a transverse section of a carpel a remarkably contracted appearance towards the commissure. This curious feature is not represented in Richard's work; the specimens there described, probably, not exhibiting perfectly ripe fruit. *J. D. H.*

Fig. 1. Flower. *f. 2.* Ripe fruit. *f. 3.* The same cut through transversely:—*magnified.*



Parkeriana.

N. O. *Orchideæ.*

TAB. CCCXIII.

SCAPHYGLOTTIS ROSEA.

Pseudobulbis elongato-cylindraceis sulcatis proliferis, foliis linearibus obtusis, floribus geminatis, labello obovato-spathulato apice recurvo trilobo lobo medio acuminato.

HAB. Demerara. *C. S. Parker, Esq.*

This is certainly very near the *S. violacea* of Dr Lindley (*Bot. Reg. t. 1901,*) a plant of the same country; but as that has also been sent us, as well as the present, in a living state, I have no hesitation in pronouncing them distinct. The flowers of this are of a much paler hue, the sepals and petals of an uniform rose colour: but the chief characteristic lies in the labellum, which is here attenuated below into a long narrow claw, while the upper half is curved down, the apex 3-lobed, the middle coming to a rather sharp point.

The habit of this genus is peculiar in its proliferous pseudobulbs: but the structure of the flower seems to me scarcely different from that of *Maxillaria*.

Fig. 1. Flower. *f. 2.* Column and lip. *f. 3.* Lip:—*magnified.*



TAB. CCCXIV.

EPIDENDRUM RIGIDUM. Jacq.

Caule erecto ancipiti-compresso, foliis distichis linear-i-oblongis coriaceis obtusis, spicæ rachi ancipiti, sepalis ovatis coriaceis, petalis oblongis membranaceis, labelli postici cordato-ovati obtusi concavi palato calloso cuniculo inflato, columna apice dentato-laciniata, bracteis foliaceis dolabriformibus carinatis ovarii longitudine.

Epidendrum. Jacq. Amer. p. 222. t. 134. Sw. Fl. Ind. Oc. 3. p. 1507. Lindl. Gen. et Sp. Orchid. p. 110.

β. labello subrotundo. *Lindl. l. c. (an species distincta?)*

HAB. West Indian islands. Our specimen here figured is from St Vincent. *Rev. L. Guilding.*—β. Mexico. (*Lindley.*)

A frequent inhabitant of Jamaica and other of the West Indian islands, and sometimes cultivated in our stoves, though possessing little beauty to recommend it.

Fig. 1. Flower. f. 2. Portion of do. f. 3. Column, from which the anther is removed. f. 4. Anther-case. f. 5. Pollen-masses:—magnified.



Parkerianæ.

N. O. Orchidææ.

TAB. CCCXV.

MAXILLARIA NANA.

Humilis, caule folioso, foliis linear-lanceolatis subfalcatis rigidis, floribus solitariis axillaribus (ochroleucis), columna superne antice bialata, labello oblongo trilobo lobo medio obtusissimo cum acuminulo, disco glandula aurantiaca.

HAB. Demerara. *C. S. Parker, Esq.*

A small species of *Maxillaria* with somewhat of a proliferous habit, and hence approaching *Scaphyglottis*: at the setting on of the leaves and of the flowers also, there are copious membranaceous, sheathing scales. The sepals and petals are erect, cream-coloured. The column is prolonged below the apex of the germen, so as with the base of the labellum to form an obtuse spur. Pollen-masses 4, unequal, attached to a large, 2-toothed, membranous gland.

Fig. 1. Flower. *f. 2.* Column and lip. *f. 3.* Pollen-masses. *f. 4.* Back view of the masses removed from the gland:—*magnified.*



Parkerianæ.

N. O. Leguminosæ.

TAB. CCCXVI.


ACACIA SETIGERA.

Glabra, ramis acute trigonis, phyllodiis fasciculatis ternis lineariibus uninerviis integerrimis apice setigeris, spicis axillaribus phyllodia superantibus, petalis 4 reflexis.

HAB. New Holland? (*C. S. Parker, Esq.*)

From the greenhouse of my friend Mr Parker, who is ignorant of its native country. It is a small shrub, remarkable for its triangular branches, and the ternate, narrow, one-nerved, setigerous phyllodia and the copious racemes of bright yellow flowers that spring from the axils of the superior leaves.

Fig. 1. Portion of a branch with leaves. *f. 2.* Single flower :
—magnified.



Parkeriana.

N. O. Orchidææ.

TAB. CCCXVII.

SCAPHYGLOTTIS? FASCICULATA.

Pseudo-bulbis fasciculatis lanceolatis sulcatis monophyllis, folio linearis retuso costato avenio, pedunculo solitario brevi unifloro ad basin folii bracteato, bracteis numerosis imbricatis, perianthio patente, sepalis ovatis concavis, petalis angustioribus oblongis, columna brevi, labello oblongo obsoletissime trilobo recurvo.

HAB. Demerara. *C. S. Parker, Esq.*

Of the Genus of this plant I am extremely doubtful. It was received in a living state, but the anther had fallen away, and no pollen-masses were to be found.

Fig. 1. Flower, side view. *f. 2.* Front view of do. (the anther having fallen away). *f. 3.* Column and labellum :—*magnified.*



TAB. CCCXVIII.

HELIPTERUM INCANUM.

Incaneo-tomentosum, caule herbaceo adscendente aut erecto simplici apice aphylo, foliis linearibus integerrimis subacutis uninerviis, superioribus distantibus multo minoribus, capitulo terminali nudo, involucri subcampanulati squamis longis stipitatis ad apicem stipitis lanato-ciliatis productis in laminam lanceolatam, achaenii glabris. *De Cand.*


Helipterum incanum. De Cand. Prodr. v. 6. p. 215.

Helichrysum incanum. Hook. Bot. Mag. t. 2881.

HAB. Interior of N. Holland. *All. Cunningham, Esq.* Frequent in Van Dieman's Land. *R. Gunn, Esq.* (Nos. 108, 124, 239, 442, 837.) *Mr Lawrence*, (n. 217).

According to our views of this species it is extremely variable, and passes into the *H. bicolorum* of Allan Cunningham and De Candolle. The stems are from 6 inches to a foot and more high: the leaves more or less narrow and more or less dense, silky or hoary, or even almost woolly; the scales of the involucre are sometimes white, occasionally tinged with brown or purple. The Genus *Helipterum* of De Candolle only differs from *Helichrysum*, in the plumose, not scabrous pappus.

Fig. 1. Inner scale of the involucre. *f. 2.* Floret. *f. 3.* Hair of the pappus:—*magnified.*



Jamesonianæ,

N. O. Musci.

TAB. CCCXIX.

DICRANUM MACRODON.

Cæspitosum, caulis erectis brevibus subramosis, foliis undique insertis erecto-patentibus subulatis paululum falcatis integerim, nervo mediocri infra apicem evanescente, seta elongata, capsula erecta angusto-pyriformi, operculo e lata basi subulato obliquo, peristomii dentibus elongatis profunde bipartitis in conum approximatis.

HAB. Pillzhum, Province of Cuença, S. America? Prof. W. Jameson?

An apology is certainly necessary for giving so very imperfect an account of this species of a Genus requiring more than an ordinary degree of care in its investigation. The fact is, the drawing was made by Dr J. D. Hooker, and the plate prepared previous to his departure on the scientific voyage in the southern hemisphere under Capt. James Ross: but the specimen and original drawing, and the notes made by him, have been mislaid, and the above is all the information we can venture to give. Our impression is that the species is one of Dr Jameson's from Pillzhum. We trust at some future time to make up for our present deficiency of knowledge relative to this Moss.

Fig. 1. Tuft: nat. size. f. 2. Leaves. f. 3. Capsule, with operculum. f. 4. Capsule, showing the peristome. f. 5. Single tooth of the peristome:—magnified.



Gunnianæ.

N. O. Compositæ.

TAB. CCCXX.

HELICHRYSUM GUNNIANUM.

Caule erecto vel subdecumbente ramoso gracili arachnoideo-tomentoso, foliis acutissimis pubescentibus subtus incanato-tomentosis, inferioribus subspathulatis superioribus sensim minoribus linearibus, ramis monocephalis flavis, involuci hemisphærici squamis obovatis uninerviis denticulatis obsolete venosis, stipitibus lanatis ad apicem longe barbatis.

HAB. Van Dieman's Land. *Mr R. Gunn.* (*n.* 502.)

A most extensive Genus, even now that *Helipterum*, DC. is excluded, and one whose species we can hardly hope to define satisfactorily without the assistance of figures. With regard to the Genus of the present plant, it is hard to say whether the hairs of the pappus should come under the denomination of scabrous or plumose, which is the only distinction between the two genera just mentioned.

Fig. 1. Scale of the involucre. *f. 2.* Floret. *f. 3.* Hair from the pappus;—*magnified.*



Jamesonianæ.

N. O. Filices.

TAB. CCCXXI.


POLYPODIUM PILIGERUM.

Fronde brevi-stipitata parce pilosa lato-lanceolata bipinnatifida, laciiniis obscure costatis primariis anguste lanceolatis secundariis ovato-oblongis integerrimis obtusis, soris solitariis globosis ad basin laciniarum.

HAB. Pillzhum, Province of Cuença. S. Am. Prof. W. Jameson.

Caudex horizontalis, brevis, squamosus, squamis densis, imbricatis, nigro-fuscescentibus, nitidis. Stipites vix unciam longi, graciles, filiformes. Frons digitalis ad spithameam, circumscriptione lato-lanceolata, profundissime pinnatifida, subrigide membranacea, pilis sparsis patentibus, praecipue ad marginem et costam, obsita; laciiniis primariis unciam sesquiunciam longis patentibus, anguste lanceolatis, pinnatifidis, segmentis ovato-oblongis, modice patentibus, integerrimis, obtusis. Sori solitarii, globosi, ad basin segmentorum seu laciniarum segmentorum.

Fig. 1. Portion of the frond seen from above. *f. 2.* Portion of the same seen from beneath, with sori:—*magnified*.



Nuttallianæ.

N. O. Rosaceæ.

TAB. CCCXXII.

CERCOCARPUS BETULÆFOLIUS. *Nutt.*

Foliis rhomboeobovatis dentatis supra glabris subtus (junioribus etiam) ad nervos solummodo adpresso pubescentibus, floribus solitariis erectis v. inclinatis.

Cercocarpus betulæfolius. Nutt. Mst. in Herb. nostr.

HAB. Santa Barbara, Upper California. *Mr Nuttall.*

The accompanying figure is taken from Mr Nuttall's specimen in my Herbarium, and exhibits all the characters I am acquainted with of this species. The fruit is quite unknown to me.



Douglasiana.

N. O. Rosaceæ.

TAB. CCCXXXIII.

CERCOCARPUS PARVIFOLIUS. Nutt.

Foliis late obovatis dentatis subtus (junioribus præcipue) pubescenti-hirsutis, floribus geminatis recurvis.

Cercocarpus parvifolius. Nutt. in Herb. nostr.—Torr. et Gr. Fl. N. Am. ined.—Hook. et Arn. Bot. of Beech. Voy. Suppl. p. 337.

HAB. California. *Douglas.* Platte river of the same country.
Nuttall.

Fig. 1. Flower. *f. 2.* Calyx laid open, to show the stamens and pistil. *f. 3, 4.* Anthers. *f. 5.* Fruit with the calyx-tube (the limb being deciduous). *f. 6.* Fruit, from which the calyx-tube is removed. *f. 7.* Fruit laid open, showing the seed:—*magnified.*





CERCOCARPUS LEDIFOLIUS. Nutt.

Ramulis villosissimis, foliis oblongo-lanceolatis marginibus revolutis utrinque subtus præcipue dense pubescenti-tomentosis, floribus aggregatis erectis, staminibus exsertis.

Cercocarpus ledifolius. Nutt. *mst.* in *Herb. nostr.*

HAB. Bear River of the Rocky Mountains, California. Mr Nuttall.

Frutex, ut videtur, parvus; ramis tortuosis, crassiusculis, vetustis cicatricatis glabris, junioribus foliosis, lanato-vilosissimis. *Folia* fasciculata, coriacea, lato-lanceolata, seu oblongo-lanceolata, utrinque sed subtus præcipue fuscescenti-tomentosa, brevissime petiolata. *Flores* fasciculati, ad apices ramulorum congesti. *Calyx* dense pubescens. *Stamina* sub-20, exserta; *filamenta* quam in aliis speciebus longiora. *Fructus* cauda elongata, sericeo-vilosissima instructus.

This very distinct species of *Cercocarpus* has only hitherto been found by Mr Nuttall, on the Bear River, one of the numerous tributaries of the Rio Sacramento which falls into the Pacific Ocean at San Francisco.

Fig. 1. Flowering, and *f. 2.* fruiting specimen: *nat. size.* *f. 3.* Flower. *f. 4.* Stamen. *f. 5.* Fruit. *f. 6.* Leaf:—*magnified.*



TAB. CCCXXV.

ACTINOLEPIS MULTICAULIS. DC.

GEN. CHAR. *Actinolepis. DC.*—*Capitulum* pluriflorum, fl. radii 3-5-ligulatis stamineis, disci tubulatis 5-dentatis bisexualibus, stylo abortivo masculis? *Invol.* ovato-oblongum bracteis foliaceis paucis cinctum, squamis paucis obtusis adpressis dorso molliter tomentosis. *Recept.* angustum nudum. *Ligule* late breves 2-3-dentate, stylo bifido exerto. *Tubuli* tubo tereti, sauce dilatata, antheris albis, stylo simplici apice capitellato aut subnullo? *Achænia* oblonga subangulata, radii pubentia pappo squamellis circ. 5 scariosis acuminatis constante superata, disci glabra calva.—*Herba annua tenella multicaulia California.* Caules tenues apice subaraneosi. Folia opposita parva late obovata, basi cuneata et integra, apice obtusissima 5-dentata. Capitula in axillis supremis approximatis solitaria subcongesta parva. Flores flavi.—An *Helenia* ex pappo radii squamellato? An *Partheniea* si discus vere masculus? *DC.*

Actinolepis multicaulis. De Cand. Prodr. v. 5. p. 656. Hook. et Arn. Bot. of Beech. Voy. p. 353.

HAB. California. *Douglas.*

By some strange error the stems are stated in Decandolle to be about “2” inches in length.

Fig. 1. Capitulum and floral leaf. f. 2. Central floret. f. 3. Radial floret and scale of the involucre. f. 4. Achénium of the ray. f. 5. Scale from the pappus:—magnified.



Douglasianæ.

N. O. Compositæ.

TAB. CCCXXVI.

MADARAGLOSSA HETEROTRICA. DC.

Pilis sparsis brevibus remotis aliis pubiformibus aliis glandula nigra terminatis subscabrida, caule ramoso, foliis linearis-oblongis obtusis integris aut vix subserratis, ligulis apice trifidis involucro duplo fere longioribus. *DC. Prodr. v. 5. p. 694.*

Layia heterotricha. Hook. et Arn. Bot. of Beech. Voy. Suppl. p. 358.

HAB. California. *Douglas.*

Plant 1-2 feet high. Flowers large. The inner scales of the involucre embrace the base of the radial floret with two wings in the lower half which lap over the ovary.

Fig. 1. Outer floret of the disk with its accompanying scale. f. 2. Floret of the ray with its scales of the involucre. f. 3. Achenium of the ray. f. 4. Hairs of the pappus:—magnified.



Galeottiana.

N. O. Piperaceæ.

TAB. CCCXXVII.

PEPEROMIA GALEOTTIANA.

Acaulis humilis, foliis longe petiolatis ovato-acuminatis glaberrimis trinerviis, petiolo gracili, scapis petiolum æquantibus, spicis elongatis gracilibus folia superantibus.

HAB. Xalapa, Mexico. *Galeotti.*

An elegant and graceful little *Peperomia* with its leaves all radical, thin and membranaceous, 3-nerved, the lateral nerves slightly branched.



Galeottiana.

N. O. Anonaceæ.


TAB. CCCXXVIII.

ANONA? BIBRACTEATA.

Foliis ovato-ellipticis subcordiacis utrinque acutis nervis subtus pubescentibus, pedunculis unifloris folio vel ramo oppositis inferne bibracteatis, bracteis ovatis membranaceis foliaceis inæqualibus, sepalis ovatis petalisque (6) æqualibus lato-lanceolatis pubescentibus.

HAB. Xalapa, Mexico. *Galeotti.*

I have seen nothing of the fruit of this plant, nor yet a flower in an entirely fit state for examination, so that I am unable to decide satisfactorily even on its Genus. It is remarkable for the large, unequal, leafy bracteas situated towards the base of the peduncle; which peduncle, as well as the floral coverings, is clothed with short down.



Galeottianæ.

N. O. Berberideæ.

TAB. CCCXXIX. CCCXXX.


BERBERIS (MAHONIA) FRAXINIFOLIA.

Inermis, foliis pinnatis foliolis 7-9 ovato-lanceolatis basi acutis coriaceo-membranaceis subsessilibus integerrimis utrinque reticulatis, racemis versus apicem ramorum folium æquantibus pendulis, pedicellis gracillimis flore triplo quadruplo longioribus ebracteolatis.

HAB. Xalapa, Mexico. *Galeotti.*

This is a shrub, or probably a small-sized tree, with large, pinnated leaves, quite destitute of spines or serratures, and with leaflets not inaptly resembling those of the common Ash. The racemes too are peculiarly long and slender, and the pedicels about an inch in length. The fruit I have not seen.

Fig. 1. Petal and Stamen. f. 2. Pistil:—*magnified.*



Galeottiana.

N. O. Ternstrœmiaceæ.

TAB. CCCXXXI.

SAURAUJA BARBIGERA.

Ramulis petiolis pedunculisque ferrugineis-pubescenti-setosis, foliis obovato-lanceolatis acuminatis basi attenuatis mucronato-serratis utrinque (sub lente) punctato-scabris subtus ad axillas nervorum dense barbato-lanatis, calycibus glabriusculis.

HAB. Xalapa, Mexico. *Galeotti*.

Frutex, ramis vetustioribus glabris. *Folia* tactu et sub lente punctis minutis elevatis scabrida. *Panicula* sublonge pedunculata, corymbosa. *Calycis sepala* subrotunda, ciliata, dorso glabriuscula. *Stamina* numerosa, ad basin petalorum, subpolyadelpha. *Filamenta* basi paululum latiora, hirsuta. *Antheræ* lineares, versatiles, loculis superne poro dehiscentes. *Ovarium* globosum, 5-loculare, loculis ad angulos interiores pluriovulatis. *Styli* 5, subtorti.

Fig. 1. Flower. *f. 2.* Stamens and petal. *f. 3.* Calyx and pistil. *f. 4.* Section of the ovary. *f. 5.* Portion of the under-side of a leaf. *f. 6.* Stamen :—*more or less magnified*.



Galeottianæ.

N. O. Piperaceæ.

TAB. CCCXXXII.

PEPEROMIA LANCIFOLIA.

Caulescens suberecta, foliis lato-lanceolatis acuminatis pinnatim nervosis basi in petiolum attenuatis, pedunculo versus apicem ramorum folio duplo breviore, spicis geminatis elongatis gracilibus.

HAB. Xalapa, Mexico. *Galeotti.*

I have in vain endeavoured to find any description which will correspond with this plant, of which the solitary specimen in my possession has the spikes geminate; but I am unable to say if that character be constant.



TAB. CCCXXXIII.

GARRYA FADYENII.

Dioica, foliis ellipticis brevissime apiculatis, spicis masculis ramosis pendulis, fœmineis simplicibus erectis, bracteis flores superantibus, masc. sepalis apice cohærentibus, fœm. stylis brevibus crassis ovario adpressis.

HAB. Woods upon the Blue Mountains, Jamaica. *Dr M'Fadyen.*

The very remarkable *Genus* to which this plant belongs, was established by Dr Lindley in 1834, on a new plant of North California, found by Mr Douglas, but discovered many years previously by Mr Menzies in his voyage with Capt. Vancouver, and existing in several Herbaria to which he liberally presented it. It was, therefore, a matter of great astonishment to me, to find the same *Genus* in a plant of Jamaica, to which Dr M'Fadyen directed my attention about four years ago, and which is here represented. Mexico, however, which may be reckoned an intermediate country, is now known, by the exertions of Mr Hartweg, to produce three other species, which are described by Mr Bentham in his excellent "*Planta Hartwegianæ.*" Mr Skinner has lately sent me a species, in fruit only, from Guatemala: but this will probably prove to be identical with one or other of Hartweg's.

Our plant turns very black in drying. The connate bracteas are longer than the flowers and somewhat leafy. The segments of the male perianths are connate with their apices. The pistils are obovate. Ovules 2, pendent from the top of the cells. Styles thick, acuminate, close pressed to the top of the germen. Berry black, glabrous. Perfect seed 1. Embryo small, placed obliquely towards the upper extremity of a hard, fleshy albumen.

Fig. 1. Male spike: *nat. size.* *f. 2.* Portion of the same. *f. 3.* Single flower. *f. 4.* The same, with the perianth forced open. *f. 5.* Female spike. *f. 6.* Pistil. *f. 7.* The same laid open. *f. 8.* Berry. *f. 9.* The same laid open. *f. 10.* Seed *cut* *through* vertically. *f. 11.* Embryo:—*magnified.*



Douglasianæ.

N. O. Compositæ.


TAB. CCCXXXIV.

HARTMANNIA? PUNGENS. Hook. et Arn.

Herbacea basi sublignosa confertim foliosa, caule parce ramoso pilis albidis setoso, foliis inferioribus pinnatifidis, lobis oblongis v. oblongo-lanceolatis spinoso-mucronatis, superioribus ac fasciculorum axillarium linearibus integris rigidis spinescentibus margine recurvis papilloso-pilosis, capitulis subsolitariis bracteatis, involuci squamis spinescentibus glandulosis ligulas radii bifidas subæquantibus, achæniis epapposis, radii gibbis areola terminali valde obliqua, receptaculo toto paleaceo, paleis lanceolato-subulatis spinescentibus. *Hook. et Arn. Bot. of Beech. Voy. Suppl. p. 357.*

HAB. California. *Douglas.*

Fig. 1. Scale of the involucre, and radial floret. *f. 2.* Side view of a radial floret. *f. 3.* Back view of do. *f. 4.* Scale of the involucre. *f. 5.* Palea and central floret. *f. 6.* Central floret:—*magnified.*



TAB. CCCXXXV.

COTULA MYRIOPHYLLOIDES. Harv.

Aquatica repens glabra, ramis erectis, foliis oppositis coadunatis fere ad basin capillaceo-multifidis, capitulis globosis, involucri foliolis 4 ovatisæqualibus simplici serie, radii flosculis foemineis apetalis achaenii stipitatis alatis, disci flosculis hermaphroditis corolla tubulosa 4-fida achaenii subsessilibus apteris.

HAB. Watery places, near Cape Town. Hon. W. H. Harvey.

A very distinct and remarkable species of *Cotula*, for a knowledge of which, as well as for the drawing here given, we are indebted to our very acute friend, the Hon. W. H. Harvey. The leaves are extremely curious, and the florets still more so. Those in the circumference of the capitulum are female, upon a distinct, fleshy, cellular stipes, destitute of corolla, and with the germen and achenium winged, 2-toothed, sometimes unequally so, at the top. Central florets nearly sessile, perfect: the achenium wingless: the corolla is rather infundibuliform, being enlarged upwards, and 4-toothed.

Fig. 1. Capitulum. *f. 2, 3.* Florets of the circumference. *f. 4.* Fruit of the same. *f. 5.* Floret of the disk or centre. *f. 6.* Stigmas. *f. 7.* Achenium of the disk. *f. 8.* Leaves:—*magnified.*



Harveyana.

N. O. Cyttineæ.

TAB. CCCXXXVI.

CYTINUS DIOICUS. Juss.

Dioicus pauciflorus, perianthio campanulato muriculato 6-lobo,
limbo erecto, ovario 6-costato.

Cytinus dioicus. Juss. in *Ann. du Mus.* v. 12. p. 443.

Phelypea sanguinea. Thunb. *Fl. Cap.* p. 2.

Hypolepis sanguinea. Pers.—Spreng. *Syst. Veg.* v. 1. p. 19.
Harv. Gen. S. Afr. Plants. p. 300.

HAB. Parasitical on the roots of *Eriocephalus racemosus*, on the
sides of the Table Mountain, facing Camps Bay, at an eleva-
tion of about 2000 feet. Hon. W. H. Harvey.

Whole plant of a clear blood-red colour. Except in the
truly dioecious habit, 6-parted perianth, and erect limb, this
species offers no character by which it can be generically separat-
ed from *Cytinus*; and as these appear, in the absence of other
characters, to be of merely specific value, I fully agree with
Jussieu and Endlicher in uniting it with that Genus. W. H. H.

TAB. CCCXXXVI. Male and young plants, *natural size*,
showing their position on the woody portion of the root of
Eriocephalus. Fig. 1. Male perianth cut open. f. 2. Staminal
column. f. 3. Female perianth. f. 4. Sepal. f. 5. Style and
stigma, the lobes of the perianth being removed. f. 6. Trans-
verse section of ovary:—all but f. 1 and 3 *magnified*.



Grahamianæ.

N. O. Proteaceæ.

TAB. CCCXXXVII.


ANADENIA MANGLESII. Grah.

Foliis cuneiformibus triplinerviis venosis utrinque glabris et subconcoloribus trifidis lobis cuneiformibus, lateralibus trifidis intermedio 3-5-fido segmentis mucronatis. *Grah.*

HAB. New Holland. Introduced to our gardens by *Mr Lowe.*
(*Graham.*)

An erect, quite glabrous shrub; with harsh, rigid leaves, and racemes that are terminal or axillary, many-flowered. Flowers small, in pairs. Perianth tetraphylous, at first opening only between the claws, cohering by the capitate apex (as at f. 1.), afterwards opening entirely. Pistil stipitate; stipes white, curved. Ovary gibbous, yellow, bearing 3 ovules. Style conical, more swollen on one side, white. Stigma conical, pink-coloured. *Graham.*

Fig. 1. Appearance of a flower on first opening. *f. 2.* Flower, fully expanded. *f. 3.* Pistil :—*magnified.*



Douglasianæ.

N. O. *Onagrarieæ.*

TAB. CCCXXXVIII.

ENOOTHERA (HOLOSTIGMA) GRACILIFLORA. *Hook. et Arn.*

Acaulis pilosa, foliis anguste linearis-spathulatis integerrimis v.
apice denticulatis, floribus sessilibus, calycis tubo elongato
filiformi folia subæquante, petalis lato-obcordatis. *Hook. et*
Arn. in Bot. of Beech. Voy. Suppl. p. 341.

HAB. California. *Douglas.*

A very decidedly marked species of an extensive genus. Root annual, long, tapering, slender. Leaves all radical, 2-4 inches in length, hoary with longish soft hairs. From the top of the somewhat fusiform ovary, arises the very long slender tube of the downy calyx, which brings the corolla nearly on a level with the apex of the leaves. The petals are yellow, green when dry.

Fig. 1. Flower:—magnified.



TAB. CCCXXXIX.


ŒNOTHERA (HOLOSTIGMA) ALYSOIDES. Hook. et Arn.

Humilis multicaulis puberula, foliis inferioribus multo majoribus oblongo-lanceolatis inæqualiter dentatis in petiolum attenuatis caule paullo brevioribus, superioribus linearibus, racemis foliosis secundis circinnatis, ovaris tenuissimis sessilibus, petalis filamenta æquantibus stylo brevioribus, capsulis contortuplicatis striatis torulosis. *Hook. et Arn. Bot. of Beech. Voy. Suppl. p. 341.*

HAB. Pine Creek, Snake Country; California. *Tolmie.*

One of the most remarkable of the *Holostigma* group of *Œnothera*. It has a slender, perpendicular, branching root, from the summit of which arise several short ascending stems, from 3 to 5 inches long, the central one, however, is erect, and always apparently floriferous from the base to the summit, the capsule at the base being almost mature, while the upper portion is only in flower. Lower leaves almost equal in length to the stems, and sometimes nearly an inch broad, while the upper ones become bracteas and are shorter than the flowers. Corolla pale yellow, retaining their colour when dry, as in *Œ. dentata*, (Cav.); they are copious, racemose, and secund, the upper portion of the raceme being recurved or circinate until the flowers expand. Petals roundish, not at all notched. The whole habit of the plant is indeed very similar to that of some *Alyssum* or *Vesicaria*.

Fig. 1. Flower and bractea. f. 2. Fruit:—magnified.



TAB. CCCXL.

FABIANA IMBRICATA.


Ramis virgatis, foliis undique imbricatis cylindraceis punctatis basi productis, floribus terminalibus solitariis in ramulis brevibus subsessilibus.

Fabiana imbricata. *Ruiz et Pav. Fl. Peruv. v. 2. p. 12. t. 122.*
Lindl. Bot. Reg. t. 59.

HAB. Chili. *Ruiz et Pavon.* Andes of Chili. *Mr Cruickshanks.* *Mr Cuming* (n. 140). *Dr Gillies.* Coast of South Chili. *Dr Eights.*

Our drawing was taken from a living specimen, raised in the Edinburgh Botanic Garden, from seeds gathered by Mr Cuming: and since our plate was finished it has appeared in the Botanical Register. Although even the coloured figure makes but little show on paper, in consequence of the pure white of the flowers, yet the number and gracefulness of these among the copious dark green foliage of the numerous branches, have a very pretty effect. The inflorescence on a short peduncle. Calyx nearly globose, shortly and bluntly 5-lobed. Corolla funnel-shaped. Limb plaited, of 5 rounded, spreading lobes. Ovary with large red fleshy scale, or hypogynous gland on each side. Ovary 2-celled, many-seeded. Fruit 2-valved. Receptacle of the fruit eventually separating from the valves and cohering at the top.

Fig. 1. Flower. *f. 2.* The same laid open. *f. 3.* Transverse section of the hypogynous glands and ovary. *f. 4.* Capsule. *f. 5.* Receptacle of the seeds removed from the capsule:—*magnified.*



Galeottiana.

N. O. Ternstroemiaceæ.

TAB. CCCXLI. CCCXLII.


SAURAUJA PEDUNCULATA.

Dioica? ramis petiolis pedunculisque nudiusculis, foliis late obovato-oblongis acutis basi obtusis v. acutis serratis utrinque laevibus, calycibus dense pubescentibus.

HAB. Xalapa, Mexico. *Galeotti.*

A much larger plant than our *S. barbigeria*, figured at tab. 331, with leaves broader, a span long, very indistinctly serrated, with longer, more glabrous peduncles, and very downy calyces. The flowers which I examined appeared to be all male, having a small abortive pistil, and very short erect styles.

Fig. 1, 2. Flowers. *f. 3.* Petal and stamens. *f. 4.* Single stamen. *f. 5.* Calyx and abortive pistil. *f. 6.* Pistil from the same. *f. 7.* Section of do.:—*magnified.*



Douglasianæ.

N. O. Composite.

TAB. CCCXLIII.

MONOLOPIA MINOR. DC.

GEN. CHAR. *Capitulum* multiflorum heterogamum, fl. radii fœmineis ligulatis interdum subbilabiatis, disci tubulosis bermaphroditis sterilibus aut masculis, tubo tenui hispidulo, fauce dilatata. *Invol.* squamæ 8-10 uniserialis ad medium concretae. *Recept.* convexum aut subconicum epaleaceum. *Antheræ* ecundatæ. *Styli* radii rami cono brevi superati. *Achania* calva glabra, radii compressa aut subtrigona, disci compressa. —Herbæ *Californicæ annuæ erectæ albo-tomentosæ*. Folia alterna oblongo-linearia integra aut parce dentata aut lobulata. Capitula ad apicem caulis aut ramorum terminalia ebracteata. Flores lutei.—Genus accedit ex invol. ad *Gamolepidem*, disco sterili ad *Steirodiscum*. *De Cand.*

M. *minor*; foliis linearibus, lobulis paucis hic inde instructis, radicalibus glabratiss., cæteris albo-tomentosis, ligulis involucre parvo longioribus. *De Cand. Prodr. v. 6. p. 74. Hook. et Arn. in Bot. of Beech. Voy. Suppl. p. 359.*

HAB. California. *Douglas.*

A Californian Genus of two species, the present and that figured in the following plate. In this, when the involucre comes to be accurately examined, it will be seen that the scales of the involucre are not in one series, neither are they so much united as in *M. major*.

Fig. 1. Involucre. f. 2. Radii floret. f. 3. Floret from the disk: magnified.



Douglasianæ.

N. O. Composite.

TAB. CCCXLIV.

MONOLOPIA MAJOR. DC.


Foliis ligulatis obtusis vix subdentatis, ligulis involucro triplo longioribus.

Monolopia major. *De Cand. Prodr.* v. 6. p. 74. *Hook. et Arn. Bot. of Beech. Voy. Suppl.* p. 359.

HAB. California. *Douglas.*

This should be considered the type of the Genus *Monolopia*; of which the character was given in our last page. The involucre is here truly formed of one series of leaflets, and there are some other peculiarities which I do not perceive in *M. minor*. The florets of the ray are furnished with a curious toothed scale, as shown at *f. 5*, and the teeth of the florets of the disk are beset externally with spreading jointed hairs. (See *f. 2, 3.*)

Fig. 1. Receptacle. *f. 2.* Floret of the disk. *f. 3.* Hair from the teeth of the corolla. *f. 4.* Floret from the ray. *f. 5.* Portion of the same, to show the scale at the top of the tube. *f. 6.* Achénium. *f. 7.* Section of the same:—magnified.



Forbesiana.

N. O. Vacciniæ.

TAB. CCCXLV.


VACCINIUM FORBESII.

Foliis semperferventibus late ovatis subobovatisve subcoriaceis obtusiusculis superne præcipue serratis subtus reticulatis, racemis axillaribus, pedicellis 2-3-bracteatis bracteis ovatis membranaceis, dentibus calycinis acutis, corollæ campanulatae 5-fidæ lobis ovatis, filamentis hirsutis, antheris linearibus muticis, stylo exerto.

HAB. Mozambique. *Forbes.*

Few Genera of plants are more universally dispersed than this, from Alpine heights on mountains to the lowest plains, from the tropics almost to arctic regions, and in both hemispheres. The plant here figured is an inhabitant of the eastern coast of Africa, in lat. 15° south. Other species are indigenous to the neighbouring island of Madagascar.

Fig. 1. Flower. f. 2. Stamen. f. 3. Pistil:—magnified.



Hartwegiana.

N. O. Euphorbiaceæ.

TAB. CCCXLVI.

EUPHORBIA SPHÆRORHIZA. *Benth.*

Glabra, radice tuberosa sphærica, caule herbaceo erecto dichotomo, foliis linearibus integerrimis utrinque angustatis, involucris solitariis breviter pedicellatis, laciniis interioribus brevisimis fimbriatis, glandulis truncatis dorso in appendiculas ovatas fimbriatas coloratas productis. *Benth.*

Euphorbia sphærorrhiza. *Benth. Pl. Hartw. p. 8.*

Hab. Zacatecas, Mexico. *Hartweg.*

An *Euphorbia* very distinct from any hitherto described, with the habit, as Mr Bentham observes, of *E. Ipecacuanha*, but very different in the involucres and in the extremely narrow leaves. Root a globose bulb, with a few fibres from below. Involucre cup-shaped, 1-2 between each upper pair of leaves, or from the axil of the upper dichotomies, hairy, with 5 glands at the margin, each spreading out into a white petaloid 2-3-cleft appendage. Inner laciniae alternating with these, short, membranous, bifid, and denticulate.

Fig. 1, 2. Involucres with male flowers. *f. 3.* Gland from the margin: *f. 4.* Inner lacinia:—*magnified.*



Smithianæ.

N. O. Orchideæ.

TAB. CCCXLVII.

EPIDENDRUM MICROBULBON.

Pseudo-bulbo parvo ovato vaginato diphylo, foliis lato-linearibus scapo brevioribus, racemo paucifloro, sepalis petalisque conformibus, labello sublibero oblongo trilobo disco lamellato lamellis undulatis, lobo intermedio subrotundo integerrimo crispato.

HAB. Oaxaca, Mexico. *Robert Smith, Esq.*

Introduced by Robert Smith, Esq., of Oaxaca, to the gardens at Woburn Abbey. It would belong to the *Encyclia* division of *Epidendrum*, or that group which has the labellum almost entirely free from cohesion with the column. Its nearest affinity is perhaps with the *Encyclia patens* (Bot. Mag. t. 3013, *Epidendrum odoratissimum*, Lindl. Bot. Reg. t. 1415), and is chiefly distinguished by its narrower sepals and petals, not at all spathulate.

Fig. 1. Column. f. 2. Labellum:—magnified.



Parkinsonianæ.

N. O. Orchideæ.

TAB. CCCXLVIII.


MAXILLARIA ANGUSTIFOLIA.

Pseudo-bulbis proliferis oblongis levibus vetustis subsulcatis, pedunculis bulbo longioribus unifloris, sepalis petalisque conformibus oblongis, labello oblongo obscure trilobo disco tuberculo oblongo lobø medio ovato.

HAB. Mexico. J. Parkinson, Esq.

Sent by Mr Parkinson to Woburn in 1838. The pseudo-bulbs rise one above another in a proliferous manner, and are more or less sheathed with membranous, brownish scales. The single-flowered peduncles spring solitary from the base of each pseudo-bulb. The perianth is of a tawny yellow colour, the labellum of a deeper tint.

Fig. 1. Column, petals, and labellum. *f. 2.* Back view; and *f. 3.* Front view of the pollen-masses:—*magnified*.



TAB. CCCXLIX.

RUBUS ROSÆFOLIUS.

Caule teretiusculo piloso, aculeis patentibus subrecurvis, foliis pinnatis pilosiusculis, pinnis lanceolatis duplicato-serratis glanduloso-punctatis, stipulis linear-setaceis, pedunculis sub-unifloris, laciinis calycinis lanceolatis longe acuminatis petalis vix longioribus, carpellis glabris numerosissimis minoribus exsiccatione lacunoso-rugosis. DC.

Rubus rosæfolius. Sm. Ic. Ined. 3. p. 60. t. 60. *De Cand. Prodr. v. 2. p. 556.*


R. coronarius; petalis numerosis calyce multo longioribus. Sims in Bot. Mag. t. 1783. *De Cand. Prodr. l. c.*

Rubus Sinensis. Hortul.

Rubus Commersonii. Poir. Dict. 6. p. 242. (*fide De Cand.*)

HAB. Mauritius. Thouin (in Smith). Molucca islands, Bojer (in Hort. Maurit.), who observes that it is naturalized in Mauritius. Table Mountain, Cape of Good Hope, Ecklon. Foot of Table Mountain. Hon. W. H. Harvey.

Smith gives this plant, on the authority of Thouin, as an inhabitant of the Mauritius. Bojer, who has so long resided in that country, tells us it is naturalized there, and comes from the Moluccas. De Candolle, who also states Mauritius to be its native country, nevertheless adduces the *R. Commersonii*, Poir. as a synonym: but that is said to have a reddish fruit the size of a raspberry, but with a much less agreeable taste and smell: from which it may be inferred that it is juicy; while the fruit of our plant is, according to Mr Harvey, as dry "as in *Potentilla*." Ecklon and Mr Harvey (and to this latter gentleman we are indebted for the drawing here figured) find the species, to all appearance wild, about Table Mountain, Cape of Good Hope, and perhaps that is really its native country.



Hartwegianæ.

N. O. Olacineæ.

TAB. CCCL.


XIMENIA PARVIFLORA. *Benth.*

Spinosa, ramis flexuosis, foliis oblongo-lanceolatis, floribus axillaribus glomeratis 4-5-meris 8-10-andris. *Benth. Plant. Hartweg. p. 7. n. 28.*

HAB. Zacatecas, Mexico. *Hartweg. (n. 28.)*

Rami angulati, rigidi. *Spinae* fere in omnibus axillis. *Flores* in racemulis axillaribus brevissimis 5-10-floris dispositi, dimidio fere minores quam in *X. Americana*. *Pedicelli* oppositi. *Bractæ* minute. *Calyx* minutus, lacinia acutis. *Petala* siccitate nigra, intus ad medium rufo-barbata, barba multo breviore quam in *X. Americana*. *Filamenta* filiformia. *Antheræ* oblongæ, filamento ipso longiores. *Stylus* brevis, conicus, stigmate simplici acutiusculo. *Ovarium* breve, carnosum, 3-4-(5?)loculare, ovulo in quoque loculo solitario pendulo. *Benth. l. c.*

Fig. 1. Flower. *f. 2.* Petal. *f. 3.* Stamen. *f. 4.* Anther cut through transversely. *f. 5.* Pistil cut through transversely: —*magnified.*



Zeyherianæ.

N. O. Scrophularinæ.

TAB. CCCLI.

HARVEYA PURPUREA. Harv. mst.


Calyce profunde 5-fido lobis lanceolatis tubi corollæ pubescen-
tis inflato-ventricosi dimidium æquantibus, corollæ limbo
planiusculo margine undulato.

Orobanche Uitenhagensis. Eckl.

Harv. Uitenhage, South Africa; Zeyher.

Our character of the genus *Harveya* is given at TAB. CXVII. of this work, together with a figure of the only species then known to us. The present is distinguished from it by the different form of the calyx, and of that of the tube of the corolla, which is, moreover, hairy or downy, and by the less crisped limb of the latter. The scales are more numerous on the scape or stem, and the colour, judging from the name, is purple, not pale rose. It is one of the many discoveries of Mr Zeyher in Uitenhage, and the drawing is made from the dried specimen of our friend Mr Harvey at the Cape.

The specific character of *H. Capensis* may stand thus:—
Calyce campanulato breviter 5-fido, tubo corollæ glabro cylindraceo superne sensim dilatato, limbi margine undulato eroso-crispato. (TAB. CXVII.)



TAB. CCCLII.

PAPPEA CAPENSIS, Eckl. et Zeyh.

GEN. CHAR. *Flores abortu diclines. Cal. inæqualiter 5-partitus. Petala 4—6, intus barbata.—MASC.: Stamina 8—10, disco annulari hypogyno inserta, filamentis villosis.—FÆM.: Ovarium 3-loculare, loculis 1-ovulatis. Stylus 1 (ex 3 coalitis.) Stigma sub-3-fidum. Carpella 3, capsulari-drupacea, basi connata, globosa, sepe 1—2 abortiva.—Arbor 15—20-pedalis. Folia obvato-oblonga, obtusa, margine revoluta, coriacea, glaberrima, ramulorum apice congesta. Racemi nudi, ramulorum apice congesti; masc. folia subæquantes; fæm. 4—6-flori brevissimi.*
—Eckl. et Zeyher.

Pappea Capensis, Eckl. et Zeyh. En Pl. Afr. Austr. p. 53; Harv. Gen. S. Afr. plants, p. 37; Arnott in Journ. Bot. 3. p. 258.

Kiggelaria integrifolia, E. M. in Dredge's herb. (excl. syn. Jacq.)

Its fruit called “*Wild Preume*,” is fleshy, and is eaten, and an oil is expressed from the seeds.—*Kiggelaria integrifolia*, Jacq., judging from the figure given by that Botanist, is totally unlike the present plant. In it the leaves are scattered over the branches, and not collected at their extremity; the branches are larger and more simple; its leaves are pubescent on both sides, and dark green on their upper surface; the racemes more lax and fewer flowered, while the flowers themselves are cernuous; the petals are large, as in *Kiggelaria*, and longer than the stamens; the plant is polyandrous, and there are two diverging styles. Moreover, it appears that *Kiggelaria integrifolia* and *Pappea* were both found by Ecklon and Zeyher, and referred the one to *Flacourtiææ*; the other, and correctly so, to *Sapindaceæ*.—ARN.

Fig. 1. Male flower, side view. *f. 2.* Ditto, front view. *f. 3.* Inner view of a petal. *f. 4.* Outer view of do. *f. 5.* Stamen. *f. 6.* Fruit, nat. size. *f. 7.* Fruit with one cell, or lobe, laid open: all but *f. 6.* magnified.



Adamiana.

N. O. Passifloretæ.

TAB. CCCLIII.


PASSIFLORA LESCHENAUTII.

Scandens, foliis semiorbicularibus truncatis tricuspidatis subtus pubescentibus eglandulosis, petiolis medio biglandulosis, cirrhis simplicibus, pedunculis geminis unifloris, calyce involucrato, petalis 5, fructu ovoideo stipitato, stipite calycis longitudine.

Passiflora Leschenaultii. *De Cand. Prodr.* 3. p. 326. *Wall.*
Cat. n. 1231. *Wight, Cat.* n. 1154. *Wight et Arn. Prodr.*
Fl. Pen. Ind. Or. 1. p. 353.

HAB. East Indies, Neelgherries; *Dr Wight.* At Mayaburan;
Sir Frederic Adam, G.C.B.

Our specimen of this Passion-flower, a Genus of very rare occurrence in the Old World, was kindly presented to me with a considerable collection of other plants from the Neelgherries, by Sir Frederic Adam, late governor of Madras, and I have thought it deserving of a figure in this little work. The colour of the flower appears to be white; the rays are long, not very crowded, and equal in length with the petals.



Tolmieana.

N. O. *Hydrophyllæ*.

TAB. CCCLIV.

EUTOCA? LUTEA.

Nana diffusa pubescenti-scabra ramosa, radice multiplici annus, foliis petiolatis subrotundo-ovobatis basi cuneatis crenatolobatis superioribus subintegerrimis, ovariis 8-ovulatis, corolla campanulata marcescente persistente (lutea) calycem paullo superante.

Eutoca? lutea. *Hook. et Arn. in Bot. of Beech. Voy. p. 373.*

HAB. Snake Fort, Snake Country, California; *Mr Tolmie.*

This is a very remarkable and a beautiful species, with bright yellow flowers (looking at the first glance like those of some cruciferous plant.) The seeds are deeply wrinkled on the back. It differs from *Eutoca* by the persistent but marcescent corolla, and from *Emmenanthe* by its habit and by the absence of the scales of the corolla.

Fig. 1. Flower. *f. 2.* Corolla laid open. *f. 3.* Pistil. *f. 4.* Fruit with persistent calyx and corolla. *f. 5.* Capsule. *f. 6.* The same burst open, showing the seeds. *f. 7, 8.* Seeds:—*magnified.* .



Tolmieanae.

N. O. *Hydrophyllæ.*

TAB. CCCLV.

EUTOCA ARETIODIES.

*Nana diffusa hispido-pilosa trichotome ramosa, radice multici-
piti annua, foliis petiolatis spathulato-lanceolatis basi longe
attenuatis integerrimis, ramulis ultimis brevibus congestis,
floribus sessilibus terminalibus et in dichotomiis, calycis pro-
funde 5-partiti lacinias anguste linearibus hispidis tubo co-
rollæ infundibuliformis (purpureæ) duplo brevioribus, ovario
multiovulato.*

Eutoca aretioides. *Hook. et Arn. in Bot. of Beech. Voy. Suppl. p.
374.*

β. perpusilla; caule subnullo paucifloro. *Hook. et Arn. l. c.*
HAB. Burnt and Malheur rivers, Snake Country, California;
communicated by *Mr Tolmie.*

A widely different species from any *Eutoca* hitherto described, and as already observed by Dr Arnott and myself, it may be found to constitute a distinct genus. The corolla is singularly long and funnel-shaped; the stamens unequal; the calyx segments almost setaceous, and persisting when the frond is ripe.

Fig. 1. Flower. *f. 2.* Pistil. *f. 3.* Stamens. *f. 4.* Fruit and persistent calyx. *f. 5.* One valve of the fruit with seeds:—*magnified.*



TAB. CCCLVI.

WIMMERIA. Cham. et Schlecht.


GEN. CHAR. *Flores hermaphroditæ, regulares, pentameri, præfloratione imbricati. Cal. parvus, obtuse lobatus. Petala* sub margine disci inserta, sessilia, subconcava, marcescentia. *Stami.* ibidem inserta, cum petalis alternantia. *Antheræ* bilobulares lateraliter et longitudinaliter dehiscentes. *Discus* crenatus, lobatus. *Germen* disco impositum, basi immersum, pyramidato-trigonum, triloculare, loculis pluri-ovulatis; ovulis basi affixis. *Style* brevis. *Stigma* trilobum, terminale. *Pericarpium* samaroides, triquetrum, tripterum, abortu uniculare, monospermum.—Arbusculæ glabrae, habitu Celastræ, foliis exstipulatis alternis integris serratis, inflorescentis cymosis axillaribus.—*Cham. et Schlecht.*

Wimmeria concolor; foliis concoloribus.—*Cham. et Schlecht. in Linnæa*, v. 6. p. 428. *Benth. Pl. Hartweg*, p. 9. n. 41.

HAB. Mexico, near Xalapa; Scheide and Deppe. Zacatecas. *Hartweg*, n. 41.

The character of this genus depends on the indehiscent, one-seeded, samaroid fruit, according to the authors of it. Mr. Benthams justly observes that the peduncles here are only from 1—3-flowered. They are more divided according to Chamisso and Schlechtendal.

Fig. 1. Flower-buds. *f. 2.* Flower. *f. 3.* Pistil and hypogynous disk. *f. 4.* Ovary cut through transversely:—*magnified.*



TAB. CCCLVII.

MOZINNA SPATHULATA. β .

GEN. CHAR. *Mozinna*. *Orteg.* *Loureira*. *Cav.* — *Flores* dioici. *Cal.* 5-partitus. *Cor.* urceolata, 5-loba. *Glandulae* 5.—
MASC. : *Stam.* 8—13, filamentis infra coalitis, 5 exteriora brevia. — *Fœm.* : *Style* bifidus. *Stigmata* 2, late emarginata, aut 4 linearia. *Capsula* calyce persistente infra cincta, 2-valvis, 2-cocca, coccis monospermis, interdum abortu 1-locularis, 1-sperma. — *Frutices succo fætido gummoso*. *Folia alterna aut fasciculata, stipulacea, integra aut rarius lobata, pilis glanduliferis interdum (sicut calyces) ciliata*. *Flores bracteati, axillares aut terminales, feminei solitarii geminative, masculi fasciculati aut corymbosi*. — *Adr. Juss.*

Mozinna spathulata; foliis cuneatis integris v. trifidis eglandulosis. *Ortega*, *Dec.* 8. *p.* 105. *t.* 13.

Loureira cuneifolia. *Cav. Ic. t. 5. p. 17. f. 429.*

β . *sessiliflora*; folijs omnibus integris, floribus sessilibus pubescentibus. (*TAB. NOSTR. CCLVII.*) *Benth. Pl. Hartweg*, *p. 9. n. 37.*

HAB. *Zacatecas, Mexico; Hartweg, (n. 37.)*

Stems very stout, and rather succulent than woody, wrinkled when dry, and clothed with a dark olive-coloured shining bark, bearing numerous warty excrescences, from whence the leaves and flowers arise. These flowers in our specimen are all male; and they, as well as the leaves, are very much smaller than in the plant of Ortega and Cavanilles.

Fig. 1. Male flower. f. 2. Glands and stamens from the same:—magnified.



TAB. CCCLVIII.

LOBELIA TRIGONA. Roxb.

Glabra, ramis diffusis erectis vel adscendentibus cauleque trigonis trialatisve, foliis ovatis nunc subcordatis in petiolum brevem basi angustatis dentatis dentibus mucronulatis, pedicellis gracilibus bractea longioribus basi bibracteolatis, tubo calycis obconico, lobis linearí-acuminatis tubum subæquantibus, corolla glabra lobis calycinis subdimidio longiore, antheris inclusis omnibus apice barbatis, capsula obovoidea.

Lobelia trigona, Roxb. *Fl. Ind.* I. p. 506; (*ed. Wall.*) II. p. 111; in *E. I. C. Mus. Tab. Pict.* 439; *De Cand. Prod.* 7. p. 359.

Lobelia trialata, Ham. in *Don. Prod. Fl. Nep.* p. 157; *De Cand. Prod.* 7. p. 360.

Lobelia stipularis, Roth. in *Roem. et Sch. Syst.* 5. p. 67.


Lobelia Heyneana, Roem. et Sch. *Syst.* 5. p. 50.

Lobelia micrantha, Hook. *Exot. Fl.* I. t. 44.

HAB. Ceylon, at an altitude of 6000 feet; *Col. Walker.* Peninsula of India; *Dr Wight.* Neelgherries; *Sir F. Adam.* Simla; *Lady Dalhousie.* Népal; *Dr Wallich.*

The form of the leaf, especially at the base, and the length of the petiole, is so variable that we have little doubt of these synonyms belonging to the same species. It is also the *L. Zeylanica* of many herbaria; but the Linnaean species of that name appears to be a Chinese plant with villous capsules, while the *Campanula ceilanica* of Seba, from whom Linnaeus borrowed the name, appears perfectly distinct from either, and, if really from Ceylōp, may be referred to *L. radicans*, Wall., although it has more the habit of some Cape species.—*ARN.*

Fig. 1. Small specimen from the Neelgherries. *Sir Fred. Adam.* *f. 2.* Portion from Dr Wallich's specimen. *f. 3.* Plant from Ceylon. *Colonel Walker: nat. size.* *f. 4.* Side view of a flower. *f. 5.* Capsule. *f. 6.* Portion of the stem:—*magnified.*



Nuttallianæ.

N. O. Cruciferæ.

TAB. CCCLIX.

ARABIS PUBERULA. *Nutt.*


Perennis subcespitosæ magis minusve pubescens pilis densis stellata, foliis integerrimis linear-lanceolatis sessilibus, siliquis planis rectis pendulis, pedicellis longitudinem sepalorum duplo æquantibus, seminibus leniter marginatis. *Nutt.*

Arabis puberula. Nutt. in Torr. et Gr. Fl. of N. Am. v. 1. p. 82.

Forests of the Blue Mountains of Oregon (Columbia R.),
N. America; *Nuttall.*

Our figure of this plant is taken from a specimen obligingly sent to us, with many other similar treasures, by Mr Nuttall, who observes that the specimens were all in fruit, that the pods are slightly pubescent, (scarcely apparent in our specimens,) and that the central nerve is obvious.

Fig. 1. Pod:—magnified.



TAB. CCCLX.

MECONELLA OREGANA. Nutt.

Meconella Oregonæ, Nutt. in Torr. et Gr. Fl. of N. Am. v. 1.
p. 64.


HAB. Open plains of the Oregon (Columbia R., N. W. Am.)
near its confluence with the Walamet; Nuttall.

"An interesting but humble plant," Messrs Torrey and Gray
observe, "which seems to stand between *Platystemon* and *Hypocoum*, two genera having apparently little relationship, yet that
ought, notwithstanding their anomalies, not to be removed from
Papaveraceæ. It agrees with the latter in its definite stamens,
and with the former in the foliage and floral envelopes, dilated
filaments, &c. The torus is somewhat like that of *Chryseis* on
a small scale."

The following is Mr Nuttall's character of this genus:—
Sepals 3, somewhat pilose. Pet. 5—6. Stamens 4—6. Filaments membranaceous, dilated* upwards; anthers very short,
the cells disjoined. Stigmas 3, (rarely 4) linear, very short,
sessile. Capsule pod-shaped, slender, 3 (rarely 4) valved: valves
flat, 1-nerved, debiscing from the apex to the base, not separating
from the placenta. Seeds numerous, smooth, and shining,
subglobose.—A very small, annual, glabrous, and somewhat
glauous herb. Stamens slender, dichotomously branched.
Radical leaves in a rosulate cluster, spatulate; caudine ones
linear or linear-spatulate, opposite; the lowermost ternate,
entire. Peduncles axillary, filiform, 1-flowered. Flower very
small, ochroleucous.

Fig. 1. Flower-bud, with calyx. *f. 2.* Expanded flower. *f. 3.*
Capsule:—*magnified.*

* In our dried specimens the dilatation of the filaments is hardly percep-



TAB. CCCLXI.

OXALIS CRATENSIS.

Herbacea erecta ramosa glabra, caule folioso, foliis sublonge petiolatis unijugis cum impari petiolulato subrhombeo-ovatis obtusis, pedunculis axillaribus folium æquantibus apice bifidis racemosis, rachibus (floribus delapsis) denticulatis, sepalis ovatis non sphacelatis, filamentis glabris, stigmatibus patentibus, staminibus brevioribus iis longioribus.

HAB. Moist cane fields about Crato, Brazil; *Mr Gardner.*

Root branching, annual. Stems 8 or 10 inches to a foot high, rounded, herbaceous, glabrous. Leaves on rather long slender petioles, bearing 3, ovato-rhomboïd, glabrous leaflets, about an inch long, the two lateral ones smaller, nearly sessile, the terminal one on a petiolule one-fourth the length of the leaflet. Peduncles axillary, about as long as the leaf, bifid at the extremity, each branch being a raceme of flowers which elongates as the flowers fall off, leaving a toothed rachis. Calyx glabrous: sepals ovate, acute, green, not sphacelate nor spotted. Petals pale pink, united by their bases. Filaments 5 long, and 5 short, united at the base into a distinct tube, glabrous. Anthers short, all alike. Germen ovato-globose. Styles spreading, the dilated stigmas protruded between the filaments, shorter than the shortest stamens.

Nearly allied to *O. Barrellieri* of Jacq. I have not Jacquin's figure to refer to. Lamarck describes that plant as fruticose, while ours is certainly annual. Sir James Smith, however, speaks of it as annual, and Barrellier's figure evidently appears to be so too, and is indeed a good representation of our plant. Our figure was taken from a living specimen which was raised from seeds sent by Mr Gardner.

Fig. 1. Flower. f. 2. Portion of the corolla. f. 3. Stamens and pistil. f. 4. Pistil:—magnified.



TAB. CCCLXII.

KANDELIA RHEDEI.

GEN. CHAR. *Kandelia*, *Wight et Arn.*—*Calyx* 5-fidus; laciniae tubo 3—4-plio longiores, lineares, patentes. *Petala* 5, membranacea, glabra, basi linearis subcanaliculata, ultra medium bifida, laciniis in fila plurima capillaria longa inæqualiter fissis. *Stamina* petalorum numero 6—8-pla; *filamenta* subulato-capillaria, calycis laciniis fere æquantia: *antheræ* oblongæ, parvæ, obtusæ, dorso paullo supra basia affixa. *Ovarium* adhærens, 1-loculare, 6-ovulatum. *Stylus* filiformis *stamina* subsuperans. *Stigma* 3-dentatum. *Fructus* oblongus, basi laciniis calycinis patentibus coronatus, tubo multo longior.—*Arbor parva*. *Folia anguste elliptico-oblonga*, *obtusa*. *Pedunculi petiolo longiores*, 2—3-chotomi, 4—9-flori. *Flores majusculi nunc rarius* 6-meri: alabastra prismaticæ. *Calyx bractea cupulata suffultus*. *Embryonis germinantis* tigellus clavato-subulatus, acutissimus.—*ARN.*


Kandelia Rhedei, *Wight et Arnott*, *Prodr. Fl. Penins. Ind. Or.* v. 1. p. 311; *Wight Cat.* n. 1042; *Arnott in Ann. of Nat. History*, v. 1. p. 365; *Griffith on Rhizophora*, p. 9.

Rhizophora Candel. *Linn. Sp. Pl.* p. 634; *De Cand. Prodr.* 3. p. 32; *Spreng. Syst. Veg.* 2. p. 235.; *Wall. Cat.* n. 4876. *Rheed. Hort. Mal.* v. 6. t. 35.

HAB. Malabar and Bengal.

Mr William Griffith of the Madras Medical service, in his memoir on *Rhizophoreæ*, approves of the genus constituted by Drs Wight and Arnott, but observes that the character of the ovary as given by them is at variance with the usual structure and with his own observations made on living plants. Dr Arnott, however, about the same time repeated his analysis and found no appearance of dissepiments. Perhaps they are extremely fugacious and disappear by the drying of the specimen.

Some apology is necessary for figuring this plant without any dissections; but the drawing was too hastily, and without my knowledge, sent to the printers and transferred to the stone; and in this kind of art it is well known that additions cannot be so easily made, after the proofs are taken off, as in copperplate engraving.



TAB. CCCLXIII.

FUMARIA MICRANTHA, *Lag.*

Foliorum segmentis anguste linearibus canaliculatis ultimis brevibus, bractea spathulato-lineari acuta pedicellum patulum subæquante, sepalis peltatis orbicularibus basi subcordatis inciso-dentatis margine undulatis dorso concavis corolla sub-duplo brevioribus ac 1½—2-plo latioribus, fructu globoso subapiculato.

F. micrantha. *Lag.* *Cat. Hort. Madr.* p. 21. *DC. Syst. Veg. II.* p. 137; *Prodr. I.* p. 130. *Arnott in Third Report Bot. Soc. Edin.* p. 104, 107.

F. calycina. *Bab. in Fourth Report Bot. Soc. Edin.* p. 34.

HAB. Spain; *Lagasca*. Montpelier; Dr Walker Arnott. Scotland; first discovered in the neighbourhood of Edinburgh, by D. Stewart, Esq., but since met with by various collectors in East Lothian, Perthshire, and Forfarshire.

We believe that no specimen named by *Lagasca* exists in any herbarium; but as the beautiful character derived from the sepals is so distinctly pointed out by that Botanist in the words "calycibus cordato-rotundatis corollæ tubo latioribus," we presumed that no doubt could exist as to the propriety of referring our plant to his. Mr Babington, however, on the authority of our friend Dr Klotzsch of Berlin, says that *Lagasca's* is different; but as no proof of this is offered, no notice given of the points in which the two differ, no description made of the specimen in the possession of Dr Klotzsch, so that one might judge how far it agrees with *Lagasca's* character, nor any mention made whence that specimen was obtained, we are still disposed to adhere to our former opinion.—*ARN.*

Fig. 1. Flower. f. 2. Fruit:—both magnified.

Fig. A, Flower (*magnified*) of *F. capreolata*, *y. Anglicæ*, Arn. in *Third Report, Bot. Soc. Ed.* p. 101, 106, or *F. capreolata* of British Botanists, which has been confounded by De Candolle with his *F. media*, (a form of *F. officinalis*), but rather resembles *F. micrantha* in its large calyx, although extremely different in other respects.



TAB. CCCLXIV.

STRYCHNOS TOXIFERA, Schomb.

(Ourari or Wourali poison of Guiana.)

Ramis scandentibus cirrhisque pilis longis patentibus rufis dense obtectis, foliis sessilibus ovali-oblongis acuminate membranaceis trinerviis utrinque pilis longis rufis hirsutis, floribus , fructibus maximis globosis. *Benth.*

Strychnos toxifera. Schomb. in Benth. Guian. Pt. in Hook. Journ. Bot. v. 3. p. 240.

HAB. British Guiana; *Schomburgk*, (n. 155.)

Every one, the readers of "Waterton's Wanderings" and those acquainted with Brande's experiments especially, is familiar with the name of Ourali or Wourali poison, whose effects are so powerful on animal life. We believe that no European has seen the plant, either in a living or dried state, till the distinguished traveller, to whom we are indebted for the specimens here figured, ascended the Rupunoony river in British Guiana. Being informed that the plant grew in the Conoco Mountains, he made a detour in order to have the gratification of seeing it; and under the guidance of some Indians, was directed to a ligneous twiner, at which they called out "Ourari," the name of the plant in Warpeshana.

"The stem," Dr Schomburgk continues, "is often more than three inches thick, and very crooked; its bark rough, and of a dark greyish colour; the branches thin, and inclined to climb; the leaves dark green, and opposite, ovate, acute, 5-nerved and veined; young branches and leaves hirsute; hairs brown, cirriferous; the cirrhi, however, not found on every branch. Fruit (see our TAB. CCCLXV.) of the size of a large apple, round, smooth, bluish green. Seeds imbedded in a pulp, consisting chiefly of a gammy matter, which is intensely bitter. We observed many heaps of the cut wood covered with palm trees, which, as the Indians told us, had been left by the Maroons. The plant grows only in two or three places, which are resorted to by the Indians from all directions, and often from a great distance."

Our TAB. CCCLXIV. represents a sterile branch: (the flowers are yet unknown to us.) The fruit is exhibited in the following plate,

Tab. COCLAT.



Schomburgkianæ.

N. O. Strychnæ.


TAB. CCCLXV.

STRYCHNOS TOXIFERA. *Schomb.*

(See the preceding description.)

This plate represents the fruit; *nat. size*, and *f.* 1. a seed, and *f.* 2. inside of one of the cotyledons, *do.*

We need scarcely inform our readers, that the well-known *nux vomica* of the shops, is the product of a plant of the same genus with the present.



TAB. CCCLXVI.

RAUWOLFIA HETEROPHYLLA. Herb. Willd.


Ramis angulatis glabris, foliis oppositis ternis quaternisque inæqualibus membranaceis ellipticis acuminatis penninerviis sub lente minute farinosis demum nudis, petiolis brevibus canaliculatis margine superne ciliato-aculeolatis, cymis pedunculatis paucifloris, baccis parvis pisiformibus.

Rauwolfia heterophylla. Willd. Herb. n. 5098. Roem. et Sch. Syst. Veget. v. 4. p. 805. Schlecht. et Cham. in Linnæa, v. 5. p. 125.

HAB. Mexico, Puente del Rey; Schiede and Deppe. Xalapa; *Galeotti.*

This plant is well described by Chamisso and Schlechtendal in the work above quoted. In my specimens the lower leaves are opposite, the rest ternate or quaternate, more or less unequal. Flowers few, small, in axillary cymes. Lobes of the small calyx rounded, imbricated. Corolla funnel-shaped, bearded within at the back of the anthers. Germin small, two-lobed, seated on an hypogynous gland. Berry or drupe, small, with two hard wrinkled nuts.

Fig. 1. Flower. *f. 2.* Portion of the corolla, to show the tuft of hairs at the back of the anther. *f. 3.* Pistil. *f. 4.* Drupe or berry. *f. 5.* The same cut open to show the nuts. *f. 6.* One of the nuts:—*magnified.*



Drummondianæ.

N. O. Leguminosæ.

TAB. CCCLXVII.


ACACIA SQUAMATA. Lindl.

Ramis phyllodiisque rectis apice recurvis eglandulosis acutis teretibus glabris, stipulis nullis, capitulis geminatis ternisve pedunculatis e squamis imbricatis deciduis erumpentibus, filamentis discoloribus.—*Lindl.*

Acacia squamata.—*Lindl. Sketch of the Botany of Swan River*, p. xv. n. 63.

HAB. Swan River Settlement, N. Holland; *James Drummond*.

A very remarkable species of the extensive leafless group of Australian *Acacia*. The *Phyllodia* resemble branches or spines, hooked at the point. In the axils of these are seen ovate scaly buds; from them the flowers proceed, and the scales fall away as the flowers expand. The anthers are yellow, the numerous filaments deep tawny.



TAB. CCCLXVIII.

VIScum FALCATUM.

Trunco tereti, ramis ancipitibus sub dichotomis dilatatis, foliis carnosis 3—5-nerviis falcatis obtusis semipedalibus, ♀ spicis axillaribus 1—3-pollicaribus sesquipollicaribusque, floribus verticillatis senis. *Cham. et Schlecht.*

Viscum falcatum. Cham. et Schlecht. in Linnæa, v. 5. p. 172.

HAB. About Xalapa; Schiede and Deppe. *Galeotti.*

Our specimens of this Mistletoe consist of undivided branches from ten inches to a foot and more long. The lower and older portion of the plant is described as terete.—The whole is glabrous and more or less glossy. Branches very much compressed and two-edged, about two lines wide in the narrowest part, becoming much dilated immediately beneath the leaves, striated longitudinally and somewhat reticulated in the dried state. Leaves opposite, articulated obliquely on the very dilated portions of the stem, 6—8 inches long, linear-lanceolate, obtuse, tapering at the base, coriaceous, 3—5-nerved, the nerves connected by oblique nervelets, often reticulated. Spikes, in my specimens solitary in the axils, sometimes two and three according to Chamisso and Schlechtendal, an inch and a half to two inches long, jointed and moniliform, that is, contracted at the joints, and there having a short, bifid, membranaceous, slightly ciliated sheath, which receives the base of the articulation above it. Articulations one-fourth of an inch long, or more, cylindrical, covered all round (not sessile) with small usually trimerous flowers which are too young to allow of my describing them.



Drummondianæ.

N. O. Leguminosæ.

TAB. CCCLXIX.

ACACIA DIPTERA.

Ramis foliaceis angustis bialatis glaucis marginatis foliorum loco
in lobis falcatis apice mucronatis inflexis productis, stipulis
nullis, pedunculis monocephalis racemosisque ancipitibus,
capitulis pedunculatis. *Lindl.*

Acacia diptera. Lindl. in Veget. of Sw. River, p. xv.

HAB. Swan River Settlement, Australia; James Drummond.

A well-defined and singular looking *Acacia* of the Phyllodium group, yet destitute of phyllodia as well as of leaves. The racemes bear copious heads of yellow flowers.



Drummondianæ.

N. O. Leguminosæ.

TAB. CCCLXX.

ACACIA INCRASSATA.

Stipulis parvis spinescentibus deciduis, phyllodiis coriaceis sparsis oblique triangularibus angulo unico (exteriore) mucronato altero glandulifero mutico nervo valido laterali margine undique incrassato, capitulis solitariis, pedunculis phyllodio longioribus.

HAB. Swan River Settlement, Australia; *James Drummond.*

A species closely allied to *A. decipiens*, especially to that state of it figured in the *Botanical Magazine*, t. 1745; but the phyllodia are more coriaceous, the nerve is thicker and more prominent, and what affords the best character, the margin is considerably elevated or thickened, especially at the truncated apex.



Galeottianæ.

N. O. Rosaceæ.

TAB. CCCLXXI.

PRUNUS (CERASUS) SAMYBOIDES.

Folis sempervirentibus papyraceis ovali-lanceolatis basi acutis
subtusque maculis duabus glandulosis venæ secundæ adjacen-
tibus notatis apice obtuse acuminatis glaberrimis integerrimis
subtus prominule penninerviis, racemis axillaribus subsessi-
libus solitariis aggregative, floribus 20—25-andris, stamini-
bus calyce duplo longioribus, drupis parvis subglobosis acuti-
usculis.—*Cham. et Schlecht.*

HAB. Mexico. Xalapa; (*Galeotti*) and at Hacienda de la
Laguna; *Schiede*.

The name is not inaptly applied, and we have little doubt of
this being the plant of Chamisso and Schlechtendal, though
that author does not notice the discoloured leaves, the underside
in our dried specimen being of a much browner and paler tint
than the upper one. We, on the other hand, have omitted to
notice in the figure of the under side of the leaves a constant
and very important character, namely, the *ustulate spot* on the
second vein from the base on each side of the costa. It comes
near the *P. Carolinianus*, but the leaves are less coriaceous and
the flowers are much smaller.

Fig. 1. Flower, with bractea. *f. 2.* Flower laid open:—
magnified.



Galeottianæ.

N. O. Leguminosæ.

TAB. CCCLXXII.

CROTALARIA BUPLEURIFOLIA.

Leviter pubescenti-pilosa, caule angulato subulato dichotome ramoso, foliis brevissime petiolatis ovato-ellipticis utrinque obtusis (lineari-lanceolatisque acutis) mucronatis, floribus subgeminis pedunculatis prope basin ramorum, bracteis subjunctis oppositis stipuliformibus decurrentibus apice hastato-acuminatis.—*Schl.*

Crotalaria bupleurifolia. *Schlecht.* in *Linnæa*, 5. p. 575. *Hook.* et *Arn. Bot. of Beech. Voy. Suppl.* p. 414.

β . foliis superioribus lineari-lanceolatis acutis. *Hook. et Arn. l. c.*
 γ . foliis omnibus lineari-lanceolatis acutis. *Hook. et Arn. l. c.*

HAB. Mexico.— α . Hacienda de Laguna; *Schiiede and Deppe.*
Xalapa; *Galeotti.* β . and γ . Between San Blas and Tepic;
Dr Sinclair.

This seems a very variable plant, and perhaps too nearly allied to *C. sagittalis* and *C. ovata* of the Southern States of North America.



TAB. CCCLXXIII.

MIMOSA FLORIBUNDA, H.B.K.

Fruticosa erecta, caule petiolisque aculeatis cum pedunculis ferrugineo-hispidis, aculeis sparsis, foliis conjugato-pinnatis pinnis supra demum glabriusculis subtus strigosis 4-foliolatis, quorum foliolis 3 semiovatis, quarto infimo interno abortivo minuto, pedunculis inferioribus axillaribus bi-ternisve petiolum æquantibus, superioribus racemoso-subpaniculatis, capitulis globosis densifloris.

Mimosa floribunda. H.B.K., Nov. Gen. Am. 6. p. 250. Willd. Sp. Pl. v. 4. p. 1031. De Cand. Prodr. v. 2. p. 426. Cham. et Schlecht. in Linnaea, v. 5. p. 591. Hook. et Arn. in Bot. of Beech. Voy., p. 287.

Mimosa Willdenovii. Poir. Suppl. v. 1. p. 50.

HAB. Near Cumana; Humboldt. Columbia; Cuming, (n. 1241).

Mexico. Tepic; Capt. Beechey. Xalapa; Schiede, *Galeotti*.

Stems woody, erect, and, as well as the petioles, prickly, (the prickles short, slightly curved and scattered,) and they the peduncles are densely clothed with short spreading rigid and fulvous hairs. Leaves conjugate and pinnate, each pinna consists of four leaflets, with small subulate stipules at the base; the upper pair large, equal, semiovate, the inferior pair extremely unequal, of which the outer one resembles the upper ones, the inner exceedingly small, all mucronate, nearly glabrous above, beneath, clothed with close-pressed rigid hairs or bristles of a tawny colour, those at the margin being peculiarly hard and rigid. Peduncles 2 or 3 from the axils of many of the leaves, as long as the petioles, monocephalous; the upper ones from the abortion of the leaves form a sort of elongated panicle or raceme. Heads globose. The numerous stamens appear to have been purple.



Galeottianæ.

N. O. Loranthæ.

TAB. CCCLXXIV.

LORANTHUS SCHIEDEANUS.

Fruticosus, caule angulato, foliis lato-lanceolatis coriaceis falcatis 3—5-nerviis longe acuminatis, paniculis axillaribus terminalibusque, pedicellis bi-ternis, calyce involucrato, corolla longissima (digitali) 4-fida, staminibus 6 longitudine corollæ stylum æquantibus. *Schlecht.*

Loranthus Schiedeanus. *Schlecht, in Linnæa, v. 5. p. 172.*

HAB. Xalapa, Mexico; *Galeotti.*

This appears to be a large growing plant, with deeply angular almost winged stems and branches: leaves 5—6 inches long, penninerved as Schlechtendal accurately describes them, and subtriplinerved. The flowers are in copious corymbose panicles, 4—5 inches long, hexandrous; the calyx, truncate and nearly entire, is included at the base within a cup-shaped involucre, resembling an outer calyx.

I am indebted to Wm. Harris, Esq. of Kingsbury, for the possession of this and many other fine Mexican plants.



TAB. CCCLXXV.


DROSERA BULBOSA.

Radice bulbosa tunicata nitida, caule simplici squamoso, foliis terminalibus obovato-cuneatis erectis (parvis), pedunculis congestis unifloris foliis subduplo longioribus.

HAB. Swan River, New Holland; *James Drummond.*

The root of this has the appearance of a true bulb, and is of the size of that of our common Crocuses, externally consisting of a number of dark brown glossy layers or coats, not tinged the paper with a purple dye as do the tuberous roots of this Genus. The stem is scarcely three inches high, rather thick, and clothed with jagged brown scales, probably the remains of former leaves. The perfect leaves are few in number, eight or ten, confined, as in the preceding species, to the apex, small, obovato-cuneate, clothed on the upper surface and at the margin with glandular hairs. From within these arise 2—8 or 10 single-flowered peduncles, about twice as long as the leaves, and as well as the calyx quite glabrous.

This comes near the D. erythrorhiza, Lindl. in Sw. R. Veget. p. xx. n. 90; but that has the flowers in a pedunculated cyme and the leaves much broader. In our present plant, however, the foliage is evidently young; in some specimens, received from Mr Drummond since the figure was made, the leaves are large and conceal more of the peduncles.



TAB. CCCLXXVI.

DROSERA MACROPHYLLA.


Radice tuberosa solida, caule elongato nudo simplici apice folioso basi dilatato et cum tubere articulato, foliis (magnis) cuneatis rosulatis vix in petiolum attenuatis apice truncatis supra marginibus glanduloso-pilosis, pedunculis aggregatis folio longioribus 2—3-floris.

Drosera macrophylla. Lindl. *Sketch of Swan River Botany*, p. xx. n. 91.

HAB. Swan River, New Holland; James Drummond.

The root of this beautiful plant is a small solitary tuber, about the size of a pea, imparting, as does the base of the stem, a beautiful purple dye to the paper on which the specimen is fastened. The stem is articulated as it were on the top of this, and, at the point of attachment, much swollen, probably about to form a new tuber for the following year: the rest of the stem, about four inches long, is as thick as a crow's quill, naked, or only marked with the scars of the fallen leaves. The present year's foliage is confined to the summit of the stem, rosulate, of 10—12 leaves: these leaves are the largest of any species with which I am acquainted, two inches long, and an inch or an inch and a-half broad in the upper part. Peduncles 2—3 flowered, and as well as the pedicels and calyx perfectly glabrous.

Perhaps in this *Drosera*, as well as in that of the preceding plate, (*D. bulbosa*,) the root, in its perfect state, is covered with concentric scales. The dye given out by the tuber is singularly beautiful.



Douglasianæ.

N. O. *Cupuliferae.*

TAB. CCCLXXVII.

QUERCUS AGRIFOLIA.

Foliis subcordato-ovatis remote spinoso-dentatis utrinque glabris perennantibus, fructibus solitariis geminisque axillaribus, cupulae hemisphaericæ basi attenuatae squamis adpressis oblongis obtusiusculis glabriusculis, glande ovato-oblonga acuta subconica cupula paullo longiore.

Q. agrifolia. *Neé* in *Ann. Sc. Natur.* iii. p. 271. *Willd.* *Sp. Pl.* 4. p. 431. *Spreng.* *Syst. Veget.* 3. p. 859. *Hook. et Arn.* *Bot. of Beech.* *Voy.* p. 391.

HAB. California; *Neé.* *D. Douglas.*

A very pretty and well-marked species of oak. It was first found by *Neé* at Monterrey, and it is said also at Nutka.



Gardnerianæ.

N. O. Gesneriaceæ.

TAB. CCCLXXVIII.

GLOXINIA SARMENTIANA. *Gardn. mst.*


Herbacea annua caulescens ramosa glabra, caule gracili, foliis oppositis petiolatis ovatis grosse serratis, pedunculo brevi axillari vel terminali solitario unifloro, calyce puberulo 5-fido.

Gloxinia Sarmentiana. *Gardn. Herb. Brasil, n. 2226.*

HAB. Abundant, but rare in flower, on shady rocks in a ravine called Buráco do Inferne, about a league from Oeiras, the capital of Piauhy, Brazil. *Mr Gardner.*

It is to be hoped Mr Gardner will be able to introduce this beautiful and showy annual alive to our gardens, where it would prove highly ornamental, its flowers being of a rich purple-blue, and very large in proportion to the size of the plant. This was first made known to Mr Gardner by Dr Casimiro José Moraes Sarmento, a young gentleman of Oeiras, to whom he was also deeply indebted for many kind attentions during his stay in that city, and whose name he wishes it to commemorate.

Fig. 1. Calyx and pistil, from which the corolla had fallen:—*magnified.*



TAB. CCCLXXIX.

TAXODIUM SEMPERVIRENS. Lamb. ?

Foliis sempervirentibus subdisticbis erecto-patentibus laxe imbricatis linearibus acutis supra canaliculatis glabris subtus glaucis subpulverulentis marginatis costa prominente, galbulis (?) junioribus squamis ovatis membranaceis fuscis obtectis.
Taxodium sempervirens. Lamb. Pin. t. 643? Hook. et Arn.

Bot. of Beech. Voy. p. 392.

Abies religiosa. Hook. et Arn. Bot. of Beech. Voy. p. 160. (excl. syn. H.B.K., and of Cham. and Schlecht.)
HAB. California; Douglas. Beechey.

I regret that I am not able to give a more complete figure of a *Taxodium*, as it is supposed to be, which Mr Douglas describes as the "great beauty of the Californian vegetation, which gives the mountains a most peculiar, I was going to say, awful appearance,—something which plainly tells that we are not in Europe. I have repeatedly measured specimens of this tree 270 feet long, and 32 feet round, at three feet above the ground. Some few I saw upwards of 300 feet high, but none in which the thickness was greater than those I have instanced."

It does not so well accord with *Taxodium sempervirens* of Lambert, as to induce me to consider it decidedly that plant; but I hope the attention of Californian travellers may in future be directed to it, and that we shall be able, on some other occasion, to represent the flowers and the fruit of this most magnificent denizen of western America.

Fig. 1. Upper, and f. 2. Underside of a leaf:—magnified.



Douglasiana.

N. O. Cupuliferae

TAB. CCCLXXX.


QUERCUS DENSIFLORA.

Foliis perennantibus coriaceis petiolatis oblongo-lanceolatis obtusis basi obtusis breviter acuminatis parallele nervosis integrerimis margine revolutis junioribus fulvo-furfuraceo-tomentosis subtus pallidioribus demum glabris, amentis masculis elongatis folia superantibus densifloris valde tomentosis nunc ad basin flores paucos foemineos instructis, fructibus sessilibus, cupula brevi hemispherica dense squamosa squamis elongato-linearibus laxis sericeis, glande ovato-globosa sericea.

Quercus densiflora. *Hook. et Arn. Bot. of Beech. Voy. Suppl. p. 391.*

HAB. California; *D. Douglas.*

This remarkable plant has very much of the general appearance of a *Castanea*; the fruit, in the only specimen we possess, being situated at the base of a male catkin of the former year, while the numerous male catkins of the present year exhibit no appearance whatever of female flowers. The leaves are from $2\frac{1}{2}$ to 3 inches long, and almost an inch broad.



Jamesonianæ.

N. O. Solanææ.

TAB. CCCLXXXI.


CESTRUM VESTITUM.

Foliis alternis oblongo-ovatis acutis petiolatis supra glabris reticulatis integerrimis subtus petiolis ramulis corymbis floribusque tomento pallide fulvo dense vestitis, cymis compositis terminibus densifloris, calycibus obconicis dentibus 5 brevibus truncatis, corolla infundibuliformi tubo gracili, filamentis inclusis basi geniculatis ovarioque pubescentibus.

HAB. Near Quito; Prof. W. Jameson.

Professor Jameson marks this as a tree: perhaps rather a large shrub. It is remarkable for the dense, pale, tawny tomentum, with which the specimens are everywhere clothed, except the upper part of the leaves.

Fig. 1. Flower. f. 2. Stamen. f. 3. Pistil:—magnified.



Douglasianæ.


N. O. Cupuliferæ.

TAB. CCCLXXXII.

QUERCUS DOUGLASII.

This I take to be the young branches with male flowers of the oak described in the next page. The shape of the leaves is the same, but these are in a nascent state, and clothed with a minute cinereous down. The catkins are axillary, drooping, of several rather closely placed flowers, which are sessile on the rachis. Perianth deeply 5-cleft, having ovate, acute, serrated and ciliated segments, with about six stamens. Filaments short. Anthers large, 2 celled.

Fig. 1. Portion of a male catkin with 2 flowers. *f. 2.* Perianth, from which the stamens are removed. *f. 3.* Front view, and *f. 4.*, back view of a stamen :—*magnified*.



Douglasianæ.

N. O. Cupuliferæ.

TAB. CCCLXXXIII.

QUERCUS DOUGLASII.

Foliis membranaceis oblongo-ovalibus basi acutis petiolatis sinuato-pinnatifidis siccitate haud nigrescentibus, supra glabris subtus puberulis, lobis brevibus acutiusculis, petiolis ramulisque junioribus dense fulvo-pubescentibus, fructibus sessilibus solitariis binisve, cupula hemisphærica dense squamosa squamis ovatis convexis in appendicem submembranaceam fulvam appressam linearem obtusam productis pubescentibus, glande ovata cupulam triplo superante obtusa cum umbone conico.

Quercus. Douglasii. *Hook. et Arn. Bot. of Beech. Voy. p. 391.*
HAB. California; *D. Douglas.*

For a figure of the younger state of the plant, see the preceding plate, (TAB. CCCLXXXII.)

Fig. 1. Portion of a cupule. *f. 2.* Inner, and *f. 3.* Outer scale of do :—*magnified.*



Parkinsonianæ.

N. O. Orchideæ.

TAB. CCCLXXXIV.


MAXILLARIA CURTIPIES.

Pseudo-bulbo elliptico compresso lèvissimo monophyllo, folio lato-lineari-lanceolato acutiusculo enervi basi complicato, scapo radicali pseudo-bulbum æquante squamis fuscis membranaceis vaginato unifloro, sepalis petalisque subconformatibus oblongis unicoloribus, labello oblongo obscure trilobo intus nitidissime purpureo-fusco disco tuberculo lato elevato.

HAB. Mexico; J. Parkinson, Esq.

Drawn from the living plant, of which the pseudo-bulbs were sent by Mr Parkinson to the Woburn collection. It is a small species, offering little beauty to recommend it for cultivation, but nevertheless deserving of being recorded and faithfully represented. It will rank near *Maxillaria rufescens*, Lindl. Bot. Reg. t. 1848. The pseudo-bulbs are small and singularly smooth on the base of the surface. The leaf somewhat coriaceous. From the base of the bulb there are, one on each side, the rudiments of new plants, clothed like the quarter part of the scape, with brown sheathing scales. Flower rather small, of a yellowish buff colour, destitute of fragrance, the outer or underside of the labellum is obscurely spotted with red; the inside, as well as the front of the column, is deep red-brown and glossy.

Fig. 1. Scape and flower. *f. 2.* Labellum:—*magnified.*



Jamesonianæ.

N. O. Malvaceæ.

TAB. CCCLXXXV.

SIDA PARNASSIÆFOLIA.

Acaulis, radice fusiformi tuberosa, foliis longe petiolatis cordatis obsolete dentatis petiolis ciliatis, scapis longitudine foliorum unifloris, calycibus 5-fidis basi bibracteolatis hirsutis.

HAB. Andes of Quito. On cold and elevated situations on the road from Guamote to Achupallas, at an elevation of about twelve thousand feet above the level of the sea. Professor W. Jameson.

This was only observed in places where the turf produced a herbage of very stunted growth. The flower is light purple occasionally white.

Fig. 1. Leaf. f. 2. Scape and calyx, with bracteas :—magnified.



Schomburgkianæ.

N. O. Filices.

TAB. CCCLXXXVI.

POLYPODIUM (EUPOLYP.) DISCOLOR.



Cæspitosum, fronde lanceolata subcoriacea pinnatifida supra viridi glabra subtus pulvereo-albida laciniis oblongis obtusis integerrimis, soris marginalibus uniserialibus.

HAB. British Guiana; *Mr Schomburgk*, n. 1031.

This is very different from any *Polypodium* with which I am acquainted, and remarkable for the underside being covered with a white pulverulent substance, which contrasts curiously with the dark and naked upper surface. The costa or midrib is slightly prominent on both sides; but the venation is altogether internal, and is quite obsolete on the surface. Sori dark brown, orbicular.

Fig. 1. Portion of a frond, seen from beneath, with sori:—*magnified.*

Zyp. CUCULACEA. HUTTELLIA



TABS. CCCLXXXVII. CCCLXXXVIII.

ALLOSORUS KARWINSKII.

Fronde supradecomposita, pinnis alternis petiolulatis sterilibus ovatis ovato-oblongisve obtusiusculis serrulatis basi truncatis terminali subinæquali, fertilibus copiose paniculatis in apice frondis linearibus acuminatis.


Allosorus Karwinskii. *Kunze in Linnaea*, v. 13. p. 138. *Benth. Pl. Hartweg*. p. 54.

Ceratodactylis osmundioides. *J. Smith in Bauer et Hook. Gen. Fil.* t. 36.

HAB. Mexico, Oaxaca; *Karwinski*. Barranca del Encarnacion, near Zimapan; *Hartweg*.

One of the most striking of Ferns that has recently been made known. Our specimens are from Mr Hartweg, of whose distributed collections it is No. 411. We gave the following brief notice of it in Mr Benthams invaluable notes on Hartweg's plants:—"Mr Lambert is probably the first botanist to whom this very remarkable plant was known. Mr John Smith considers it the type of a new genus, and has described it as such in the *Genera Filicum*; but I fear it cannot be separated from that division of the old genus *Allosorus*, to which Presl has referred it, and it does seem to connect *Cryptogramma*, Br. with *Allosorus*, much more so than I had any idea of till I saw this noble species. As in *Osmunda*, the upper part of the frond is converted, as it were, into a panicle of fructification."

Fig. 1. Portion of a fertile pinnule. *f. 2.* Small portion of do., the margin laid open to show the sori on the nerves. *f. 3.* *4.* Sporangia. *f. 5.* Sporules:—*magnified*.



TAB. CCCLXXXIX.


DROSCRA STOLONIFERA. Endl.

Radice tuberosa, foliis radicalibus spathulatis rosulatis, stoloni-
bus e basi procumbente adscendentibus, foliis suborbiculatis
petiolatis quaternim verticillatis scapum racemoso-panicula-
tum æquantibus, calycibus glabris. *Endlich. En. Pl. Nov.
Holl. p. 5.*

HAB. Swan River; *Hugel. Mr James Drummond.*

Endlicher appears only to have seen this curious *Drosera* des-
titute of the tuber at the base of the root. This is as large as a
good-sized nut, and stains the paper of the herbarium with a
most beautiful purple dye. Professor Lindley observes of the
Droseraceous plants of the Swan River settlement, that "they
appear likely to be in some cases of commercial value as dyers'
plants. Every part of *D. gigantea* stains paper of a brilliant
deep purple, and when fragments are treated with ammonia they
yield a clear yellow. The bulbs of *D. erythrorhiza* and *stoloni-
fera* possess the same property; in these there is a deep scarlet
powder secreted by the scales of the bulbs, which is instantly
dissolved in ammonia, forming at first an orange-coloured fluid
of great richness, but it soon changes to the rich purple above
mentioned, which is more like the colour obtained from Archil
than any thing else to which I can compare it. Possibly these
bulbs are what Dr Milligan speaks of under the name of 'boom,'
which he says are scarlet roots, not unlike in shape or size to
tulip roots. 'They roast them in the ashes, and then pound
them between two flat stones, rubbing the latter with a ball of
earth to prevent the root adhering to it: when thus prepared
they are mucilaginous, and of a glossy black colour; they may
be considered the bread of the natives who live near the coast.'
If so, they may be easily enough obtained for the purpose of
exportation, and may assist the poorer settlers in turning to ac-
count the produce of their land."

196 CCCP



TAB. CCCXC.

POLYPODIUM (EUPOLYP.) HARTWEGIANUM.

Fronde tenerima ubique stipiteque pubescentibus profunde pin-natifida (inferne pinnata,) pinnis oppositis lacinii subalternis approximatis horizontalibus e basi lata decurrente lanceolatis acutiusculis subangulato-dentatis, pinnis deflexis, infirmis (duabus paribus) basi superiore auriculatis, auriculis transversim truncatis, soris (in omnibus pinnis laciinisque) uniserialibus ovalibus oblique transversis.


*Polypodium (Eupolyp.) Hartwegianum. Hook. in Benth. Plant.
Hartweg. p. 55.*

HAB. Mexico, on the mountain Sumata, at an elevation of 9500 feet; *Hartweg. (n. 415.)*

Candex repens, squamosus. Stipes 3-pollicaris, gracilis, fusco-viridis, hinc canaliculatus. Frons pedalis, tenera, flaccida, laciinis 2—2½ uncias longis. Venæ bis furcatæ, nigricantes, venula infima superiore apice sorisera, reliquæ ante marginem evanescentes, apicibus clavatis.

Fig. 1. Portion of a fertile pinna, with sori:—magnified.

Lindsaea curvula



TABS. CCCXCI. CCCXCII.

POLYPODIUM (EUPOLYP.) SUBPETIOLATUM.

Frondes ovato-lanceolatae pinnatae, rachibus costa venisque subtus pubescenti-hirsutis, pinnis alternis remotis lanceolato-acuminatis undique serrulatis subcoriaceo-membranaceis, inferioribus basi oblique truncatis brevissime petiolatis petiolis superne alatis, superioribus basi oblique cuneatis sessilibus, supremis basi decurrentibus adnatis, soris rotundatis uniserialibus.

Polyodium (Eupolyp.) subpetiolatum. Hook. in Benth. Pl. Hartweg. p. 54.

HAB. Regla, Mexico; *Hartweg.*

Caudex repens, crassitie pennæ anserinæ, squamosus. Stipes 4—5-pollicaris. Frons pedalis. Venæ obliquæ, 2—3—4-furcatae, venula infima superne sorifera, reliquæ fere ad marginem attingentes, apicibus clavatis.

Fig. 1. Portion of a fertile pinna with sori:—*magnified.*